

LAMP-2 siRNA (h): sc-29390

BACKGROUND

Lysosome-associated membrane proteins (LAMP) are glycosylated type I membrane proteins that play a role in the biogenesis of the pigment melanin. LAMP-1 (also designated CD107A) and LAMP-2 (also designated CD107B) are involved in a variety of functions, including cellular adhesion, and are thought to participate in the process of tumor invasion and metastasis. Newly synthesized LAMP-1 and LAMP-2 proteins are sorted at the *trans* Golgi network and are transported intracellularly via a pathway that is distinct from the clathrin-coated vesicles used for the mannose-6 phosphate receptor. LAMP-1 is expressed on the surface of thrombin-activated but not resting platelets, and it is thought to be involved in the adhesive, prothrombic properties of these cells. Both LAMP-1 and LAMP-2 are involved in maintaining lysosome acidity and protecting the lysosomal membranes from autodigestion, and their expression is increased in patients with lysosomal storage disorders.

REFERENCES

1. Febbraio, M. and Silverstein, R.L. 1990. Identification and characterization of LAMP-1 as an activation-dependent platelet surface glycoprotein. *J. Biol. Chem.* 265: 18531-18537.
2. Salopek, T.G. and Jimbow, K. 1996. Induction of melanogenesis during the various melanoma growth phases and the role of tyrosinase, lysosome-associated membrane proteins, and p90 calnexin in the melanogenesis cascade. *J. Investig. Dermatol. Symp. Proc.* 1: 195-202.

CHROMOSOMAL LOCATION

Genetic locus: LAMP2 (human) mapping to Xq24.

PRODUCT

LAMP-2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LAMP-2 shRNA Plasmid (h): sc-29390-SH and LAMP-2 shRNA (h) Lentiviral Particles: sc-29390-V as alternate gene silencing products.

For independent verification of LAMP-2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29390A, sc-29390B and sc-29390C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LAMP-2 siRNA (h) is recommended for the inhibition of LAMP-2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.


GENE EXPRESSION MONITORING

LAMP-2 (H4B4): sc-18822 is recommended as a control antibody for monitoring of LAMP-2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LAMP-2 gene expression knockdown using RT-PCR Primer: LAMP-2 (h)-PR: sc-29390-PR (20 μ l, 528 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

DATA



LAMP-2 siRNA (h): sc-29390. Immunofluorescence staining of methanol-fixed, control HeLa (A) and LAMP-2 siRNA silenced HeLa (B) cells showing diminished cytoplasmic staining in the siRNA silenced cells. Cells probed with LAMP-2 (H4B4): sc-18822.

SELECT PRODUCT CITATIONS

1. Wong, J., et al. 2008. Autophagosome supports coxsackievirus B3 replication in host cells. *J. Virol.* 82: 9143-9153.
2. Hahm, E.R., et al. 2020. The role of lysosome-associated membrane protein 2 in prostate cancer chemopreventive mechanisms of sulforaphane. *Cancer Prev. Res.* 13: 661-672.
3. Fan, Y., et al. 2021. Acetylation-dependent regulation of TPD52 isoform 1 modulates chaperone-mediated autophagy in prostate cancer. *Autophagy* 17: 4386-4400.

RESEARCH USE

For research use only, not for use in diagnostic procedures.