

Stat3 siRNA (h): sc-29493

BACKGROUND

Membrane receptor signaling by various ligands, including interferons and growth hormones such as EGF, induces activation of Jak kinases which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- α and form a heterodimer which is part of the ISGF3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 β appears to be activated by both while Stat3 α is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways.

CHROMOSOMAL LOCATION

Genetic locus: STAT3 (human) mapping to 17q21.2.

PRODUCT

Stat3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Stat3 shRNA Plasmid (h): sc-29493-SH and Stat3 shRNA (h) Lentiviral Particles: sc-29493-V as alternate gene silencing products.

For independent verification of Stat3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29493A, sc-29493B and sc-29493C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Stat3 siRNA (h) is recommended for the inhibition of Stat3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Stat3 (F-2): sc-8019 is recommended as a control antibody for monitoring of Stat3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Stat3 gene expression knockdown using RT-PCR Primer: Stat3 (h)-PR: sc-29493-PR (20 μ l, 413 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Dasgupta, P., et al. 2006. Nicotine inhibits apoptosis induced by chemotherapeutic drugs by up-regulating XIAP and survivin. *Proc. Natl. Acad. Sci. USA* 103: 6332-6337.
- Dabir, S., et al. 2011. Identification of Stat3-independent regulatory effects for protein inhibitor of activated Stat3 by binding to novel transcription factors. *Cancer Biol. Ther.* 12: 139-151.
- Kang, S.H., et al. 2012. Icariside II induces apoptosis in U937 acute myeloid leukemia cells: role of inactivation of Stat3-related signaling. *PLoS ONE* 7: e28706.
- Xu, Y., et al. 2013. Epstein-Barr virus encoded LMP1 regulates cyclin D1 promoter activity by nuclear EGFR and Stat3 in CNE1 cells. *J. Exp. Clin. Cancer Res.* 32: 90.
- Santarelli, R., et al. 2014. Stat3 activation by KSHV correlates with IL-10, IL-6 and IL-23 release and an autophagic block in dendritic cells. *Sci. Rep.* 4: 4241.
- Yang, Y., et al. 2015. FAS signaling promotes gastric cancer metastasis through Stat3-dependent upregulation of Fascin. *PLoS ONE* 10: e0125132.
- Amara, S., et al. 2016. NFAT5/Stat3 interaction mediates synergism of high salt with IL-17 towards induction of VEGF-A expression in breast cancer cells. *Oncol. Lett.* 12: 933-943.
- Joy, M., et al. 2017. The myocardin-related transcription factor MKL co-regulates the cellular levels of two profilin isoforms. *J. Biol. Chem.* 292: 11777-11791.
- Peplowski, M.A., et al. 2018. Interferon γ decreases intestinal epithelial aquaporin 3 expression through downregulation of constitutive transcription. *J. Mol. Med.* 96: 1081-1093.
- Wang, H.Q., et al. 2019. Increased autocrine interleukin-6 production is significantly associated with worse clinical outcome in patients with chronic lymphocytic leukemia. *J. Cell. Physiol.* 234: 13994-14006.
- Xiong, Y., et al. 2020. ZYZ-803, a novel hydrogen sulfide-nitric oxide conjugated donor, promotes angiogenesis via cross-talk between Stat3 and CaMKII. *Acta Pharmacol. Sin.* 41: 218-228.

RESEARCH USE

For research use only, not for use in diagnostic procedures.