

TFIID siRNA (h): sc-29503

BACKGROUND

In eukaryotic systems, initiation of transcription from protein-coding genes is a complex process requiring RNA polymerase II and broad families of auxiliary transcription factors. Such factors can be divided into two major functional classes: the basal factors that are required for transcription of all Pol II genes, including TFIIA, TFIIB, TFIID, TFIIIE, TFIIF and TFIIF; and sequence-specific factors that regulate gene expression. The basal transcription factors and Pol II form a specific multiprotein complex near the transcription start site by interacting with core promoter elements such as the TATA box generally located 25-30 base pairs upstream of the transcription start site. Binding of TFIID to the TATA element initiates assembly of the other factors into a pre-initiation complex. The TATA-binding subunit of TFIID (designated TFIIDt or TBP) from higher eukaryotes contains a highly conserved 180 amino acid C-terminal domain.

REFERENCES

1. Maldonado, E., et al. 1990. Factors involved in specific transcription by mammalian RNA polymerase II: role of transcription factors IIA, IID, and IIB during formation of a transcription-competent complex. *Mol. Cell. Biol.* 10: 6335-6347.
2. Peterson, M.G., et al. 1991. Structure and functional properties of human general transcription factor IIE. *Nature* 354: 369-373.
3. Lee, D.K., et al. 1992. TFIIA induces conformational changes in TFIID via interactions with the basic repeat. *Mol. Cell. Biol.* 12: 5189-5196.
4. Takada, R., et al. 1992. Identification of human TFIID components and direct interaction between a 250 kDa polypeptide and the TATA box-binding protein (TFIIDt). *Proc. Natl. Acad. Sci. USA* 89: 11809-11813.
5. Huisinga, K.L. and Pugh, B.F. 2007. A TATA binding protein regulatory network that governs transcription complex assembly. *Genome Biol.* 8: R46.
6. Demeny, M.A., et al. 2007. Identification of a small TAF complex and its role in the assembly of TAF-containing complexes. *PLoS ONE* 2: e316.
7. Bhattacharya, S., et al. 2007. Structural analysis and dimerization potential of the human TAF5 subunit of TFIID. *Proc. Natl. Acad. Sci. USA* 104: 1189-1194.
8. Holloway, K., et al. 2007. Action of transcription factors in the control of transferrin receptor expression in human brain endothelium. *J. Mol. Biol.* 365: 1271-1284.
9. Romier, C., et al. 2007. Crystal structure, biochemical and genetic characterization of yeast and *E. cuniculi* TAFII5 N-terminal domain: implications for TFIID assembly. *J. Mol. Biol.* 368: 1292-1306.

CHROMOSOMAL LOCATION

Genetic locus: TBP (human) mapping to 6q27.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TFIID siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TFIID shRNA Plasmid (h): sc-29503-SH and TFIID shRNA (h) Lentiviral Particles: sc-29503-V as alternate gene silencing products.

For independent verification of TFIID (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29503A, sc-29503B and sc-29503C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TFIID siRNA (h) is recommended for the inhibition of TFIID expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TFIID (TBP) (58C9): sc-421 is recommended as a control antibody for monitoring of TFIID gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TFIID gene expression knockdown using RT-PCR Primer: TFIID (h)-PR: sc-29503-PR (20 μ l, 465 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.