goat anti-mouse IgG_{2a}-HRP: sc-2970



The Power to Overtin

BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. We offer horseradish peroxidase (HRP) and alkaline phosphatase (AP) conjugated isotype-specific secondary antibodies for detection of mouse $IgG_1,IgG_{2a},IgG_{2b},IgG_3,IgM$ and IgA in Western Blot and ELISA. Pre-adsorbed HRP and AP conjugated isotype-specific secondary antibodies are also available and are recommended for use with immunoglobulin-rich samples.

SOURCE

goat anti-mouse $\lg G_{2a}$ -HRP is a pre-adsorbed, affinity purified secondary antibody raised in goat against mouse $\lg G_{2a}$ and conjugated to HRP (horse-radish peroxidase).

PRODUCT

Each vial contains 200 μg goat adsorbed IgG_{2a} in 0.5 ml of 1X PBS containing 40% glycerol.

APPLICATIONS

goat anti-mouse $\lg G_{2a}$ -HRP is recommended for detection of mouse $\lg G_{2a}$ by Western Blotting (starting dilution: 1:1000, dilution range 1:1,000-1:10,000; optimal dilution to be determined by titration) and immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400).

RECOMMENDED SUPPORT PRODUCTS

- UltraCruz™ Tissue Culture Dish, 100 mm polystyrene dish: sc-200286
- UltraCruz™ Cell Scrapers, 25 cm, sterile, 100 per case: sc-213229
- RIPA Lysis Buffer, 50 ml, cell lysis buffer with protease inhibitors: sc-24948
- CompleteTM Protease Inhibitor Cocktail Tablet, 20 tablets: sc-291300
- Electrophoresis Sample Buffer, 2X, 25 ml, reducing buffer: sc-24945
- UltraCruz™ PVDF Transfer membrane, 0.45 μm, 30 cm x 3 m roll: sc-3723
- UltraCruz™ Nitrocellulose Pure Transfer Membrane, 0.22 µm, 30 cm x 3 m roll: sc-3718
- Cruz Blot-A: sc-3901 (Western Blotting membrane with human cell line extracts from 10 different cell types)
- Running Buffer, 10X, 1 L, TRIS-Glycine WB running buffer, pH 8.3: sc-24949
- Towbin, with SDS, 10X, 1 L, WB transfer buffer pH 8.3: sc-24954
- Bovine Serum Albumin (BSA), 100 g, blocking/incubation agent: sc-2323
- TBS Blotto A, lyophilized powder in single-use bottle: sc-2333
- Western Blotting Luminol Reagent, for 2,000 cm² membrane area: sc-2048
- UltraCruz™ Electrophoresis Cell: sc-201625 : runs up to 10 or 15 sample by SDS – PAGE protein electrophoresis
- UltraCruz™ Autoradiography Film, Blue, 8 x 1, 100 sheets: sc-201697
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973

SELECT PRODUCT CITATIONS

- Xie, L., et al. 2005. cDNA expression array analysis of gene expression in human hepatocarcinoma Hep3B cells induced by BNIPL-1. Acta Biochim. Biophys. Sin. 37: 618-624.
- Aimo, L., et al. 2010. Gestational zinc deficiency affects the regulation of transcription factors AP-1, NFκB and NFAT in fetal brain. J. Nutr. Biochem. 21: 1069-1075.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com