# chicken anti-goat IgG-FITC: sc-2988



The Power to Overtin

## **BACKGROUND**

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG or against antibody fragments. Santa Cruz Biotechnology offers an extensive selection of secondary antibodies optimized for immunohistochemistry and flow cytometry, and are labeled with either biotin, FITC (fluorescein isothiocyanate), Texas Red<sup>®</sup>, TRITC (tetramethyl rhodamine iso-thiocyanate), PE (phycoerythrin), PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex with cyanin-5.5). Immunohistochemistry and flow cytometry secondary antibodies are specific for commonly used primary antibody species, including goat, rabbit, mouse and rat.

# **SOURCE**

chicken anti-goat IgG-FITC is a pre-adsorbed, affinity purified secondary antibody raised in chicken against goat IgG and conjugated to FITC (fluorescein isothiocyanate).

## **PRODUCT**

Each vial contains 200  $\mu$ g chicken IgG (pre-adsorbed with mouse and human IgG) in 0.5 ml of either PBS containing 0.02% sodium azide (for IF), or PBS containing 0.1% gel and 0.1% sodium azide (for FCM).

## **APPLICATIONS**

chicken anti-goat IgG-FITC is recommended for detection of goat IgG by immunofluorescence staining (starting dilution: 1:100, dilution range: 1:100-1:400), immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400) and flow cytometry (0.5-1  $\mu$ g per 1 x 10<sup>6</sup> cells).

# **RECOMMENDED SUPPORT PRODUCTS**

#### A. TISSUE CULTURE CELLS

- □ CrystalCruz<sup>TM</sup> Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- $_{\scriptscriptstyle \rm L}$  CrystalCruz  $^{\! TM}$  Micro Slides 75 x 25 mm; 72 frosted sides: sc-24976
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- <sup>n</sup> Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336

#### **B. FROZEN TISSUE SECTIONS**

- Drgano/Limonene Mount, non-toxic alternative to Permount, 100 ml: sc-45087
- □ UltraCruz™ Mounting Medium, aqueous-based, 10 ml: sc-24941
- n ImmunoHistoMount, aqueous-based mounting medium, 30 ml: sc-45086
- <sup>n</sup> Immuno *In Situ* Mount, for use with *in situ* hybridization, 30 ml: sc-45088

#### C. FORMALIN-FIXED, PARAFFIN-EMBEDDED TISSUE SECTIONS


- Paraffin, for the preparation of tissue samples for staining, 500 g: sc-286633
- <sup>n</sup> Xylenes, mixed isomers with ethylbenzene, 500 ml: sc-237422
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973

Texas Red<sup>®</sup> is a registered trademark of Molecular Probes (6/02).


#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **DATA**



chicken anti-goat IgG-FITC: sc-2988. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization. Antibody tested: pan-TBC1D3 (T-16): sc-161990



chicken anti-goat IgG-FITC: sc-2988. Lamin B (M-20): sc-6217. Indirect, intracellular FCM analysis of fixed and permeabilized Jurkat cells stained with Lamin B (M-20), followed by FITC-conjugated chicken anti-goat IgG: sc-2988. Black line histogram represents the isotype control, normal goat IgG: sc-3887.

## **SELECT PRODUCT CITATIONS**

- Quintana, L., et al. 2009. Early tissue patterning recreated by mouse embryonic fibroblasts in a three-dimensional environment. Tissue Eng. Part A 15: 45-54.
- 2. Mladinic, M., et al. 2010. Developmental changes of gene expression after spinal cord injury in neonatal opossums. Brain Res. 1363: 20-39.
- Enunlu, I., et al. 2011. α-class prefoldin protein UXT is a novel interacting partner of amyotrophic lateral sclerosis 2 (Als2) protein. Biochem. Biophys. Res. Commun. 413: 471-475.
- 4. Rennier, K., et al. 2012. Shear stress regulates expression of death-associated protein kinase in suppressing TNF $\alpha$ -induced endothelial apoptosis. J. Cell. Physiol. 227: 2398-2411.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com