

# chicken anti-rabbit IgG-FITC: sc-2990

## BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG or against antibody fragments. Santa Cruz Biotechnology offers an extensive selection of secondary antibodies optimized for immunohistochemistry and flow cytometry, and are labeled with either biotin, FITC (fluorescein isothiocyanate), Texas Red<sup>®</sup>, TRITC (tetramethyl rhodamine isothiocyanate), PE (phycoerythrin), PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex with cyanin-5.5). Immunohistochemistry and flow cytometry secondary antibodies are specific for commonly used primary antibody species, including goat, rabbit, mouse and rat.

## SOURCE

chicken anti-rabbit IgG-FITC is a pre-adsorbed, affinity purified secondary antibody raised in chicken against rabbit IgG and conjugated to FITC (fluorescein isothiocyanate).

## PRODUCT

Each vial contains 200 µg chicken IgG (pre-adsorbed with mouse and human IgG) in 0.5 ml of either PBS containing 0.02% sodium azide (for IF), or PBS containing 0.1% gel and 0.1% sodium azide (for FCM).

## APPLICATIONS

chicken anti-rabbit IgG-FITC is recommended for detection of rabbit IgG by immunofluorescence staining (starting dilution: 1:100, dilution range: 1:100-1:400), immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400) and flow cytometry (0.5-1 µg per 1 x 10<sup>6</sup> cells).

## RECOMMENDED SUPPORT PRODUCTS

### A. TISSUE CULTURE CELLS

- CrystalCruz<sup>™</sup> Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- CrystalCruz<sup>™</sup> Micro Slides 75 x 25 mm; 72 frosted sides: sc-24976
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336

### B. FROZEN TISSUE SECTIONS

- Organo/Limonene Mount, non-toxic alternative to Permount, 100 ml: sc-45087
- UltraCruz<sup>™</sup> Mounting Medium, aqueous-based, 10 ml: sc-24941
- ImmunoHistoMount, aqueous-based mounting medium, 30 ml: sc-45086
- Immuno In Situ Mount, for use with in situ hybridization, 30 ml: sc-45088

### C. FORMALIN-FIXED, PARAFFIN-EMBEDDED TISSUE SECTIONS


- Paraffin, for the preparation of tissue samples for staining, 500 g: sc-286633
- Xylenes, mixed isomers with ethylbenzene, 500 ml: sc-237422
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973

Texas Red<sup>®</sup> is a registered trademark of Molecular Probes (6/02).


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



chicken anti-rabbit IgG-FITC: sc-2990. Indirect, intracellular FCM analysis of fixed and permeabilized Jurkat cells stained with G<sub>p</sub> (T-20), followed by FITC-conjugated chicken anti-rabbit IgG: sc-2990. Black line histogram represents the isotype control, normal rabbit IgG: sc-3888. Antibody tested: G<sub>p</sub> (T-20): sc-378.



chicken anti-rabbit IgG-FITC: sc-2990. Indirect, intracellular FCM analysis of fixed and permeabilized Jurkat cells stained with E2F-4 (A-20), followed by FITC-conjugated chicken anti-rabbit IgG: sc-2990. Black line histogram represents the isotype control, normal rabbit IgG: sc-3888. Antibody tested: E2F-4 (A-20): sc-1082.

## SELECT PRODUCT CITATIONS

- Runyan, C.E., et al. 2005. The role of internalization in transforming growth factor  $\beta$ 1-induced Smad2 association with Smad anchor for receptor activation (SARA) and Smad2-dependent signaling in human mesangial cells. *J. Biol. Chem.* 280: 8300-8308.
- Park, C.J., et al. 2010. Postnatal changes in the expression of Claudin-11 in the testes and excurrent ducts of the domestic rabbit (*Oryctolagus cuniculus domesticus*). *J. Androl.* 32: 295-306.
- Gye, M.C., et al. 2011. Expression of coxsackievirus and adenovirus receptor isoforms in developing mouse bladder uroepithelium. *Urology* 77: 1009.e9-1009.e18.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.