

# CD19 siRNA (h): sc-29968

## BACKGROUND

CD19 is a transmembrane glycoprotein that contains two extracellular immunoglobulin-like domains. CD19 is selectively expressed on the cell surface of B-lymphocytes, where it activates intracellular signaling cascades involving both Ras and phosphatidylinositol 3-kinase pathways. Activation of CD19 results in cross-linking of the membrane protein immunoglobulin chains and the subsequent association with Src family protein tyrosine kinases (PTK). Expression of CD19 is continuous throughout B-cell development and through terminal differentiation of B-cells into plasma cells. CD19 forms functional complexes with B-lymphocyte surface proteins, including Integrin  $\beta 1$ , CD21 and CD81, which are involved in regulating B-cell development.

## REFERENCES

1. Pezutto A., et al. 1987. CD19 monoclonal antibody HD37 inhibits anti-immunoglobulin-induced B cell activation and proliferation. *J. Immunol.* 138: 2793-2799.
2. Tedder, T.F., et al. 1989. Isolation of cDNAs encoding the CD19 antigen of human and mouse B lymphocytes. A new member of the immunoglobulin superfamily. *J. Immunol.* 143: 712-717.
3. Bregni, M., et al. 1989. B-cell restricted saporin immunotoxins: activity against B-cell lines and chronic lymphocytic leukemia cells. *Blood* 73: 753-762.
4. Zhou, L.J., et al. 1992. Structure of the genes encoding the CD19 antigen of human and mouse B lymphocytes. *Immunogenetics* 35: 102-111.
5. Carter, R.H. and Fearon, D.T. 1992. CD19: lowering the threshold for antigen receptor stimulation of B lymphocytes. *Science* 256:105-107.

## CHROMOSOMAL LOCATION

Genetic locus: CD19 (human) mapping to 16p11.2.

## PRODUCT

CD19 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CD19 shRNA Plasmid (h): sc-29968-SH and CD19 shRNA (h) Lentiviral Particles: sc-29968-V as alternate gene silencing products.

For independent verification of CD19 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29968A, sc-29968B and sc-29968C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}$  C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}$  C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

CD19 siRNA (h) is recommended for the inhibition of CD19 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

CD19 (B-1): sc-390244 is recommended as a control antibody for monitoring of CD19 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CD19 gene expression knockdown using RT-PCR Primer: CD19 (h)-PR: sc-29968-PR (20  $\mu$ l, 533 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$  C and the extension temperature should be 68-72 $^{\circ}$  C.

## SELECT PRODUCT CITATIONS

1. Deaglio, S, et al. 2007. CD38/CD19: a lipid raft-dependent signaling complex in human B cells. *Blood* 109: 5390-5398.
2. Kim, A., et al. 2015. Inhibition of Lyn is a promising treatment for mantle cell lymphoma with bortezomib resistance. *Oncotarget* 6: 38225-38238.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.