

Sprouty 3 (H-95): sc-30050

BACKGROUND

Members of the Sprouty family (Sprouty 1-4) are inducible negative regulators of growth factors that act through tyrosine kinase receptors. Mammalian Sprouty homologs share a well-conserved cysteine-rich carboxy-terminal domain with their *Drosophila* counterparts. Sprouty proteins are cytoplasmic in unstimulated cells, but in cells stimulated by growth factors they anchor to the plasma membrane by palmitoylation. Sprouty 1 and 2 associate with caveolin-1 in perinuclear and vesicular structures and are phosphorylated on serine residues. Sprouty 2 can associate with c-Cbl, a downregulator of RTK signaling, and inhibit the activities of several growth factors. Unlike the widely expressed Sprouty members 1, 2 and 4, Sprouty 3 expression is restricted to adult brain and testis. Sprouty 4 is a target of the WNT/ β -catenin signaling pathway in progenitor cells. In conclusion, members of Sprouty inhibit FGF and VEGF-mediated cell proliferation, suggesting that they may regulate angiogenesis in normal and disease processes.

REFERENCES

- Lim, J., et al. 2000. Sprouty proteins are targeted to membrane ruffles upon growth factor receptor tyrosine kinase activation. Identification of a novel translocation domain. *J. Biol. Chem.* 275: 32837-32845.
- Impagnatiello, M.A., et al. 2001. Mammalian sprouty 1 and 2 are membrane-anchored phosphoprotein inhibitors of growth factor signaling in endothelial cells. *J. Cell Biol.* 152: 1087-1098.
- Ozaki, K., et al. 2001. Erk pathway positively regulates the expression of sprouty genes. *Biochem. Biophys. Res. Commun.* 285: 1084-1088.
- Mailleux, A.A., et al. 2001. Evidence that SPROUTY2 functions as an inhibitor of mouse embryonic lung growth and morphogenesis. *Mech. Dev.* 102: 81-94.
- Lee, S.H., et al. 2001. Inhibition of angiogenesis by a mouse sprouty protein. *J. Biol. Chem.* 276: 4128-4133.

CHROMOSOMAL LOCATION

Genetic locus: Spry3 (mouse) mapping to X A1.1.

SOURCE

Sprouty 3 (H-95) is a rabbit polyclonal antibody raised against amino acids 1-95 mapping at the N-terminus of Sprouty 3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Sprouty 3 (H-95) is recommended for detection of Sprouty 3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Sprouty 3 (H-95) is also recommended for detection of Sprouty 3 in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for Sprouty 3 siRNA (h): sc-41039, Sprouty 3 siRNA (m): sc-41040, Sprouty 3 shRNA Plasmid (h): sc-41039-SH, Sprouty 3 shRNA Plasmid (m): sc-41040-SH, Sprouty 3 shRNA (h) Lentiviral Particles: sc-41039-V and Sprouty 3 shRNA (m) Lentiviral Particles: sc-41040-V.

Molecular Weight (predicted) of Sprouty 3: 31 kDa.

Molecular Weight (observed) of Sprouty 3: 36-43 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or JEG-3 whole cell lysate: sc-364255.

DATA



Sprouty 3 (H-95): sc-30050. Western blot analysis of Sprouty 3 expression in HeLa (A) and JEG-3 (B) whole cell lysates.

SELECT PRODUCT CITATIONS

- Rozen, E.J., et al. 2009. Loss of Sprouty1 rescues renal agenesis caused by Ret mutation. *J. Am. Soc. Nephrol.* 20: 255-259.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **Sprouty 3 (C-2): sc-374593**, our highly recommended monoclonal alternative to Sprouty 3 (H-95).