

X11 β (M-220): sc-30135

BACKGROUND

The β -Amyloid precursor protein (β -APP) is a major constituent of the amyloid deposits in patients with Alzheimer's disease. The β -Amyloid precursor is known to interact with several proteins, including X11 and the G heterotrimeric protein APP-BP1. The neuronal, transmembrane protein X11 is known to bind to the β -Amyloid precursor protein via a phosphotyrosine binding (PTB) domain, reducing the secretion of cellular β -APP and slowing β -APP processing pathways. X11 binds specifically to the YENPTY motif, which is involved in the internalization of β -APP. Multiple splice variants of X11 have been identified, including X11 α (also designated Mint 1), X11 β (Mint 2) and X11 γ (Mint 3).

REFERENCES

1. Borg, J.P., et al. 1996. The phosphotyrosine interaction domains of X11 and FE65 bind to distinct sites on the YENPTY motif of amyloid precursor protein. *Mol. Cell. Biol.* 16: 6229-6241.
2. Okamoto, M., et al. 1997. Mints, Munc18-interacting proteins in synaptic vesicle exocytosis. *J. Biol. Chem.* 272: 31459-31464.
3. Zhang, Z., et al. 1997. Sequence-specific recognition of the internalization motif of the Alzheimer's amyloid precursor protein by the X11 PTB domain. *EMBO J.* 16: 6141-6150.
4. Russo, T., et al. 1998. Fe65 and the protein network centered around the cytosolic domain of the Alzheimer's β -amyloid precursor protein. *FEBS Lett.* 434: 1-7.
5. Borg, J.P., et al. 1998. The X11 α protein slows cellular amyloid precursor protein processing and reduces A β 40 and A β 42 secretion. *J. Biol. Chem.* 273: 14761-14766.
6. Sastre, M., et al. 1998. X11 interaction with β -amyloid precursor protein modulates its cellular stabilization and reduces amyloid β -protein secretion. *J. Biol. Chem.* 273: 22351-22357.

CHROMOSOMAL LOCATION

Genetic locus: APBA2 (human) mapping to 15q13.1; Apba2 (mouse) mapping to 7 C.

SOURCE

X11 β (M-220) is a rabbit polyclonal antibody raised against amino acids 1-220 mapping at the N-terminus of X11 β of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4 $^{\circ}$ C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

X11 β (M-220) is recommended for detection of X11 β of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for X11 β siRNA (h): sc-36849, X11 β siRNA (m): sc-36850, X11 β shRNA Plasmid (h): sc-36849-SH, X11 β shRNA Plasmid (m): sc-36850-SH, X11 β shRNA (h) Lentiviral Particles: sc-36849-V and X11 β shRNA (m) Lentiviral Particles: sc-36850-V.


Molecular Weight of X11 β : 135 kDa.

Positive Controls: SK-N-SH cell lysate: sc-2410, human brain hippocampus extract: sc-364375 or mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz MarkerTM compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz MarkerTM Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruzTM Mounting Medium: sc-24941.

DATA



X11 β (M-220): sc-30135. Western blot analysis of X11 β expression in SK-N-SH whole cell lysate (A) and human hippocampus (B) and mouse brain (C) tissue extracts.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **X11 β (B-5): sc-377060** or **X11 β (E-4): sc-376510**, our highly recommended monoclonal alternatives to X11 β (M-220).