

NOR-1 (H-180): sc-30154

BACKGROUND

Nur77 (also designated NGFI-B), Nurr1 (Nur-related factor 1) and NOR-1 (neuron-derived orphan receptor-1) constitute the NGFI-B subfamily within the nuclear receptor superfamily. Ligands for these proteins have not been identified, and, therefore, they are designated "orphan nuclear receptors." Genes of the NGFI-B subfamily are classified as immediate-early genes, which are induced rapidly, but transiently, in response to a variety of stimuli. They have been implicated in cell proliferation, differentiation and apoptosis. The human NOR-1 gene maps to chromosome 9q and encodes a protein which is expressed in heart, skeletal muscle, thymus and spleen as well as in brain, where it is developmentally regulated. Therefore, NOR-1 may be involved in regulating neural differentiation. The NOR-1 gene also undergoes chromosomal translocation with the EWS gene to produce a protein thought to affect pre-mRNA splicing.

CHROMOSOMAL LOCATION

Genetic locus: NR4A3 (human) mapping to 9q22.33; Nr4a3 (mouse) mapping to 4 B1.

SOURCE

NOR-1 (H-180) is a rabbit polyclonal antibody raised against amino acids 111-200 mapping within an internal region of NOR-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-30154 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

NOR-1 (H-180) is recommended for detection of NOR-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NOR-1 siRNA (h): sc-38842 and NOR-1 siRNA (m): sc-38843; and as shRNA Plasmid control antibody for NOR-1 shRNA Plasmid (h): sc-38842-SH and NOR-1 shRNA Plasmid (m): sc-38843-SH.

NOR-1 (H-180) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of NOR-1: 68 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, Hep G2 cell lysate: sc-2227 or NOR-1 (m): 293 Lysate: sc-179018.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



NOR-1 (H-180): sc-30154. Western blot analysis of NOR-1 expression in non-transfected: sc-110760 (A) and mouse NOR-1 transfected: sc-179018 (B) 293 whole cell lysates.



NOR-1 (H-180): sc-30154. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear and cytoplasmic localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **NOR-1 (H-7): sc-393902** or **NOR-1 (F-10): sc-393903**, our highly recommended monoclonal alternatives to NOR-1 (H-180).