

Shank 2 (H-150): sc-30192

BACKGROUND

SH3 and multiple Ankyrin repeat domains 1-3 (Shank 1-3) of the Shank/ProSAP family are molecular scaffolds in the postsynaptic density (PSD). The PSD is an electron-dense structure underneath the postsynaptic plasma membrane of excitatory synapses that anchors and clusters glutamate receptors opposite to the presynaptic neurotransmitter release site. Shank proteins contain PDZ modular domains that coordinate the synaptic localization of ion channels, receptors, signaling enzymes and cell adhesion molecules. The PDZ domain mediates protein-protein interactions via the recognition of a conserved sequence motif at the C-terminus of their target protein(s). Shank recruits β PIX and PAK to spines to regulate postsynaptic structure and interacts with NMDA receptor and metabotropic glutamate receptor complexes. Transcript splice variation in the Shank family influences the spectrum of Shank-interacting proteins in the PSDs of adult and developing brain to ensure normal development.

REFERENCES

- Lim, S., et al. 1999. Characterization of the Shank family of synaptic proteins. Multiple genes, alternative splicing and differential expression in brain and development. *J. Biol. Chem.* 274: 29510-29518.
- Sheng, M., et al. 2000. The Shank family of scaffold proteins. *J. Cell Sci.* 113: 1851-1856.
- Tobaben, S., et al. 2000. The G protein-coupled receptor CL1 interacts directly with proteins of the Shank family. *J. Biol. Chem.* 275: 36204-36210.
- Sala, C., et al. 2001. Regulation of dendritic spine morphology and synaptic function by Shank and Homer. *Neuron* 31: 115-130.
- Boeckers, T.M., et al. 2002. ProSAP/Shank proteins—a family of higher order organizing molecules of the postsynaptic density with an emerging role in human neurological disease. *J. Neurochem.* 81: 903-910.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604999. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: SHANK2 (human) mapping to 11q13.3; Shank2 (mouse) mapping to 7 F5.

SOURCE

Shank 2 (H-150) is a rabbit polyclonal antibody raised against amino acids 996-1110 mapping near the C-terminus of Shank 2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Shank 2 (H-150) is recommended for detection of Shank 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Shank 2 siRNA (h): sc-42198, Shank 2 siRNA (m): sc-42199, Shank 2 shRNA Plasmid (h): sc-42198-SH, Shank 2 shRNA Plasmid (m): sc-42199-SH, Shank 2 shRNA (h) Lentiviral Particles: sc-42198-V and Shank 2 shRNA (m) Lentiviral Particles: sc-42199-V.


Molecular Weight of Shank 2 doublet: 165/180 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Shank 2 (H-150): sc-30192. Western blot analysis of Shank 2 expression in mouse brain tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Shank 2 (A-11): sc-271834** or **Shank 2 (B-4): sc-365121**, our highly recommended monoclonal alternatives to Shank 2 (H-150).