

Aph-1 (N-20): sc-30240

BACKGROUND

Four proteins comprise the γ -secretase complex: presenilin, nicastrin, Aph-1 and PEN-2. Together, these proteins mediate cell surface signaling pathways for a variety of type I membrane proteins, notably β -Amyloid precursor protein, a protein implicated in the development of Alzheimer's disease, via intramembrane proteolysis. The proteins assemble into a proteolytically active complex in the Golgi/*trans*-Golgi network (TGN) compartments. Assembly leads to autocleavage of presenilin into two subunits to create the active site of γ -secretase, an important step in understanding the mechanisms involved in the etiology and possible treatment of Alzheimer's disease.

REFERENCES

- Kimberly, W.T., et al. 2003. Identity and function of γ -secretase. *J. Neurosci. Res.* 74: 353-360.
- Baulac, S., et al. 2003. Functional γ -secretase complex assembly in Golgi/*trans*-Golgi network: interactions among presenilin, nicastrin, Aph-1, PEN-2, and γ -secretase substrates. *Neurobiol. Dis.* 14: 194-204.
- Wolfe, M.S. 2003. γ -secretase — intramembrane protease with a complex. *Sci. Aging Knowledge Environ.* 11: 7.
- Fortna, R.R., et al. 2004. Membrane topology and nicastrin-enhanced endoproteolysis of Aph-1, a component of the γ -secretase complex. *J. Biol. Chem.* 279: 3685-3693.
- Shirovani, K., et al. 2004. Identification of distinct γ -secretase complexes with different Aph-1 variants. *J. Biol. Chem.* 279: 41340-41345.
- Hansson, E.M., et al. 2005. Aph-1 interacts at the cell surface with proteins in the active γ -secretase complex and membrane-tethered Notch. *J. Neurochem.* 92: 1010-1020.
- Ma, G., et al. 2005. Aph-1a is the principal mammalian Aph-1 isoform present in γ -secretase complexes during embryonic development. *J. Neurosci.* 25: 192-198.

CHROMOSOMAL LOCATION

Genetic locus: APH1A (human) mapping to 1q21.2; Aph1a (mouse) mapping to 3 F2.1.

SOURCE

Aph-1 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Aph-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30240 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Aph-1 (N-20) is recommended for detection of Aph-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Aph-1 (N-20) is also recommended for detection of Aph-1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Aph-1 siRNA (h): sc-105081, Aph-1 siRNA (m): sc-141150, Aph-1 shRNA Plasmid (h): sc-105081-SH, Aph-1 shRNA Plasmid (m): sc-141150-SH, Aph-1 shRNA (h) Lentiviral Particles: sc-105081-V and Aph-1 shRNA (m) Lentiviral Particles: sc-141150-V.


Molecular Weight of Aph-1: 18 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or NTERA-2 cl.D1 whole cell lysate: sc-364181.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Aph-1 (N-20): sc-30240. Western blot analysis of Aph-1 expression in HeLa (A), Jurkat (B), NTERA-2 cl.D1 (C) and JAR (D) whole cell lysates.

SELECT PRODUCT CITATIONS

- Yan, R., et al. 2013. Presenilins are novel substrates for TRAF6-mediated ubiquitination. *Cell. Signal.* 25: 1769-1779.

RESEARCH USE

For research use only, not for use in diagnostic procedures.