

eIF3 β (E-15): sc-30250

BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. Eukaryotic initiation factors (eIFs) are utilized in a sequence of reactions that lead to 80S ribosomal assembly and, ultimately, translation. The eukaryotic initiation factor-3 (eIF3) scaffolding structure is the largest of the eIF complexes and includes eIF3 α , eIF3 β , eIF3 δ , eIF3 γ , eIF3 η , eIF3 ϵ , eIF3 θ and eIF3 ζ , all of which function to control the assembly of the 40S ribosomal subunit. Association of eIF3 proteins with the 40S ribosomal subunit stabilizes eIF2-GTP-Met-tRNA^{iMet} complex association and mRNA binding, and promotes dissociation of 80S ribosomes into 40S and 60S subunits, thereby promoting the assembly of the pre-initiation complex. Overexpression of eIF3 proteins is common in several cancers, suggesting a role for eIF3 proteins in tumorigenesis.

CHROMOSOMAL LOCATION

Genetic locus: EIF3I (human) mapping to 1p35.1; Eif3i (mouse) mapping to 4 D2.2.

SOURCE

eIF3 β (E-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of eIF3 β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30250 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

eIF3 β (E-15) is recommended for detection of eIF3 β of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

eIF3 β (E-15) is also recommended for detection of eIF3 β in additional species, including canine and bovine.

Suitable for use as control antibody for eIF3 β siRNA (h): sc-60080, eIF3 β siRNA (m): sc-60081, eIF3 β shRNA Plasmid (h): sc-60080-SH, eIF3 β shRNA Plasmid (m): sc-60081-SH, eIF3 β shRNA (h) Lentiviral Particles: sc-60080-V and eIF3 β shRNA (m) Lentiviral Particles: sc-60081-V.


Molecular Weight of eIF3 β : 36 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, SJRH30 cell lysate: sc-2287 or LADMAC whole cell lysate: sc-364189.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



eIF3 β (E-15): sc-30250. Western blot analysis of eIF3 β expression in HeLa (A), SJRH30 (B) and LADMAC (C) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Poleskaya, A., et al. 2007. Lin-28 binds IGF-2 mRNA and participates in skeletal myogenesis by increasing translation efficiency. *Genes Dev.* 21: 1125-1138.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



MONOS
Satisfaction
Guaranteed

Try **eIF3 β (A-7): sc-374156** or **eIF3 β (B-6): sc-271539**, our highly recommended monoclonal alternatives to eIF3 β (E-15).