

# Lamin B2 (A-14): sc-30267

## BACKGROUND

A unique family of cysteine proteases has been described that differs in sequence, structure and substrate specificity from any previously described protease family. This family, termed Ced-3/ICE, is comprised of ICE, CPP32, ICH-1/Nedd-2, Tx, Mch2, Mch3 (ICE-LAP3 or CMH-1), Mch4, and ICE-LAP6. Ced-3/ICE family members function as key components of the apoptotic machinery and act to destroy specific target proteins which are critical to cellular longevity. Nuclear lamins are critical to maintaining the integrity of the nuclear envelope and cellular morphology. The nuclear Lamin A is cleaved by Mch2, but not CPP32. Nuclear Lamin B is fragmented as a consequence of apoptosis by an unidentified member of the ICE family. Lamin C is a splice variant of Lamin A, differing only at the carboxy terminus. Lamins A and C are identical for the first 566 amino acids, with Lamin C differing only in 6 unique carboxy-terminal amino acids.

## REFERENCES

- McKeon, F.D., et al. 1986. Homologies in both primary and secondary structure between nuclear envelope and intermediate filament proteins. *Nature* 319: 463-468.
- Fisher, D.Z., et al. 1986. cDNA sequencing of nuclear lamins A and C reveals primary and secondary structure homology to inter-mediate filament proteins. *Proc. Natl. Acad. Sci. USA* 83: 6450-6454.
- Moir, R.D., et al. 1995. The dynamic properties and possible functions of nuclear Lamins. *Int. Rev. Cytol.* 162B: 141-182.
- Fernandes-Alnemri, T.F., et al. 1996. *In vitro* activation of CPP32 and Mch3 by Mch4, a novel human apoptotic cysteine protease containing two FADD-like domains. *Proc. Natl. Acad. Sci. USA* 93: 7464-7469.
- Takahashi, A., et al. 1996. Cleavage of Lamin A by Mch2 a but not CPP32: multiple interleukin 1  $\beta$ -converting enzyme-related proteases with distinct substrate recognition properties are active in apoptosis. *Proc. Natl. Acad. Sci. USA* 93: 8395-8400.
- Duan, H., et al. 1996. ICE-LAP3, a novel mammalian homologue of the *Caenorhabditis elegans* cell death protein Ced-3 is activated during FAS- and tumor necrosis factor-induced apoptosis. *J. Biol. Chem.* 271: 1621-1625.
- Duan, H., et al. 1996. ICE-LAP6, a novel member of the ICE/Ced-3 gene family, is activated by the cytotoxic T cell protease granzyme B. *J. Biol. Chem.* 271: 16720-16724.
- Rao, L., et al. 1996. Lamin proteolysis facilitates nuclear events during apoptosis. *J. Cell Biol.* 135: 1441-1455.

## CHROMOSOMAL LOCATION

Genetic locus: LMNB2 (human) mapping to 19p13.3; Lmn2 (mouse) mapping to 10 C.

## SOURCE

Lamin B2 (A-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Lamin B2 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30267 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Lamin B2 (A-14) is recommended for detection of Lamin B2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Lamin B2 (A-14) is also recommended for detection of Lamin B2 in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for Lamin B2 siRNA (h): sc-61885, Lamin B2 siRNA (m): sc-61886, Lamin B2 shRNA Plasmid (h): sc-61885-SH, Lamin B2 shRNA Plasmid (m): sc-61886-SH, Lamin B2 shRNA (h) Lentiviral Particles: sc-61885-V and Lamin B2 shRNA (m) Lentiviral Particles: sc-61886-V.

Molecular Weight of Lamin B2: 67 kDa.

Positive Controls: U-2 OS cell lysate: sc-2295, HL-60 whole cell lysate: sc-2209 or NTERA-2 cl.D1 whole cell lysate: sc-364181.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS  
 Satisfation  
 Guaranteed

Try **Lamin B2 (F-8): sc-377379** or **Lamin B2 (2Q1130): sc-71484**, our highly recommended monoclonal alternatives to Lamin B2 (A-14).