

Nanog (N-17): sc-30331

BACKGROUND

Nanog (from "Tir Na Nog", the mythologic Celtic land of the ever young) is a divergent homeodomain protein that directs pluripotency and differentiation of undifferentiated embryonic stem cells. Nanog mRNA is present in pluripotent mouse and human cell lines and absent from differentiated cells. Human Nanog protein shares 52% overall amino acid identity with the mouse protein and 85% identity in the homeodomain. Human Nanog maps to gene locus 12p13.31, whereas mouse Nanog maps to gene loci 6 F2. Murine embryonic Nanog expression is detected in the inner cell mass of the blastocyst. High levels of human Nanog expression have been detected by Northern analysis in the undifferentiated NTERA-2 cl.D1 embryonal carcinoma cell line.

SOURCE

Nanog (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Nanog of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30331 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Nanog (N-17) is recommended for detection of Nanog of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); also recommended for detection of NanogP1 and NanogP8.

Molecular Weight of Nanog: 40 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



Nanog (N-17): sc-30331. Western blot analysis of human recombinant Nanog.

SELECT PRODUCT CITATIONS

- Eckert, D., et al. 2008. Tcam-2 but not JKT-1 cells resemble seminoma in cell culture. *Cell Tissue Res.* 331: 529-538.
- Drago-Ferrante, R., et al. 2008. Low doses of paclitaxel potently induce apoptosis in human retinoblastoma Y79 cells by up-regulating E2F1. *Int. J. Oncol.* 33: 677-687.
- Nettersheim, D., et al. 2011. NANOG promoter methylation and expression correlation during normal and malignant human germ cell development. *Epigenetics* 6: 114-122.
- Larijani, M.R., et al. 2011. Long-term maintenance of undifferentiated human embryonic and induced pluripotent stem cells in suspension. *Stem Cells Dev.* 20: 1911-1923.
- Gu, T.T., et al. 2012. Cytoplasmic NANOG-positive stromal cells promote human cervical cancer progression. *Am. J. Pathol.* 181: 652-661.
- Gazouli, M., et al. 2012. OCT4 spliced variant OCT4B1 is expressed in human colorectal cancer. *Mol. Carcinog.* 51: 165-173.
- Weltner, J., et al. 2012. Induced pluripotent stem cell clones reprogrammed via recombinant adeno-associated virus-mediated transduction contain integrated vector sequences. *J. Virol.* 86: 4463-4467.
- Marlicz, W., et al. 2012. Various types of stem cells, including a population of very small embryonic-like stem cells, are mobilized into peripheral blood in patients with Crohn's disease. *Inflamm. Bowel Dis.* 18: 1711-1722.
- Yao, X., et al. 2013. Vascular endothelial growth factor receptor 2 (VEGFR-2) plays a key role in vasculogenic mimicry formation, neovascularization and tumor initiation by Glioma stem-like cells. *PLoS ONE* 8: e57188.
- Aksoy, I., et al. 2013. Sox transcription factors require selective interactions with Oct4 and specific transactivation functions to mediate reprogramming. *Stem cells* 31: 2632-2646.
- Prado, A.A., et al. 2015. Characterization of mesenchymal stem cells derived from the equine synovial fluid and membrane. *BMC Vet. Res.* 11: 281.