

LTBP-4 (N-14): sc-30360

BACKGROUND

LTBP-4 (latent TGF β -binding protein 4) is a structural component of connective tissue microfibrils which acts as a local regulator of TGF β tissue deposition and signaling. LTBP-4 exists in at least four different forms, due to alternative splicing at the amino-terminus and at the central epidermal growth factor repeat domain. LTBP-4 mRNA is present in heart, aorta, uterus and small intestine. The gene encoding human LTBP-4 localizes to chromosomal position 19q13.2.

REFERENCES

- Giltay, R., et al. 1997. Sequence and expression of a novel member (LTBP-4) of the family of latent transforming growth factor β binding proteins. *FEBS Lett.* 411: 164-168.
- Saharinen, J., et al. 1998. Identification and characterization of a new latent transforming growth factor β -binding protein, LTBP-4. *J. Biol. Chem.* 273: 18459-18469.
- Koli, K., et al. 2001. Novel non-TGF β -binding splice variant of LTBP-4 in human cells and tissues provides means to decrease TGF β deposition. *J. Cell Sci.* 114: 2869-2878.
- Mangasser-Stephan, K., et al. 2001. Expression of isoforms and splice variants of the latent transforming growth factor β binding protein (LTBP) in cultured human liver myofibroblasts. *Liver* 21: 105-113.
- Penttinen, C., et al. 2002. Secretion of human latent TGF β -binding protein-3 (LTBP-3) is dependent on co-expression of TGF β . *J. Cell Sci.* 115: 3457-3468.
- Sterner-Kock, A., et al. 2002. Disruption of the gene encoding the latent transforming growth factor β binding protein 4 (LTBP-4) causes abnormal lung development, cardiomyopathy, and colorectal cancer. *Genes Dev.* 17: 2264-2273.
- Isogai, Z., et al. 2003. Latent transforming growth factor β -binding protein 1 interacts with fibrillin and is a microfibril-associated protein. *J. Biol. Chem.* 278: 2750-2757.
- Koli, K., et al. 2004. Disruption of LTBP-4 function reduces TGF β activation and enhances BMP-4 signaling in the lung. *J. Cell Biol.* 167: 123-133.
- Chen, Y., et al. 2005. Amino acid requirements for formation of the TGF β -latent TGF β binding protein complexes. *J. Mol. Biol.* 345: 175-186.

CHROMOSOMAL LOCATION

Genetic locus: LTBP4 (human) mapping to 19q13.2.

SOURCE

LTBP-4 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of LTBP-4 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30360 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LTBP-4 (N-14) is recommended for detection of the long form of LTBP-4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LTBP-4 siRNA (h): sc-45861, LTBP-4 shRNA Plasmid (h): sc-45861-SH and LTBP-4 shRNA (h) Lentiviral Particles: sc-45861-V.


Molecular Weight of LTBP-4: 200-300 kDa.

Positive Controls: A549 cell lysate: sc-2413 or WI-38 whole cell lysate: sc-364260.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



LTBP-4 (N-14): sc-30360. Immunoperoxidase staining of formalin fixed, paraffin-embedded human premenopausal uterus tissue showing cytoplasmic staining of glandular cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.