

GP-39 (S-18): sc-30465

BACKGROUND

Human cartilage glycoprotein 39 (GP-39), also known as YKL-40, is a glycoprotein secreted by articular chondrocytes, synoviocytes and macrophages. Serum and synovial fluid GP-39 levels are elevated in inflammatory diseases and correlate with the degree of joint destruction in rheumatoid arthritis. GP-39 is expressed in articular chondrocytes, synovial cells as well as in liver but is undetectable in muscle tissues, lung, pancreas, mononuclear cells and fibroblasts. GP-39 is a candidate autoantigen in rheumatoid arthritis and is important in the capacity of cells to respond to and cope with changes in their environment.

REFERENCES

1. Hakala, B.E., et al. 1993. Human cartilage GP-39, a major secretory product of articular chondrocytes and synovial cells, is a mammalian member of a chitinase protein family. *J. Biol. Chem.* 268: 25803-25810.
2. Liu, H.W., et al. 2000. GP-83 and GP-39, two glycoproteins secreted by human epididymis are conjugated to spermatozoa during maturation. *Mol. Hum. Reprod.* 6: 422-428.
3. De Ceuninck, F., et al. 2001. YKL-40 (cartilage gp-39) induces proliferative events in cultured chondrocytes and synoviocytes and increases glycosaminoglycan synthesis in chondrocytes. *Biochem. Biophys. Res. Commun.* 285: 926-931.
4. Tsuji, T., et al. 2002. Analysis of chondrex (YKL-40, HC gp-39) in the cerebrospinal fluid of patients with spine disease. *Spine* 27: 732-735.
5. Recklies, A.D., et al. 2002. The chitinase 3-like protein human cartilage glycoprotein 39 (HC-gp39) stimulates proliferation of human connective-tissue cells and activates both extracellular signal-regulated kinase- and protein kinase B-mediated signalling pathways. *Biochem. J.* 365: 119-126.

CHROMOSOMAL LOCATION

Genetic locus: CHI3L1 (human) mapping to 1q32.1; Chi3l1 (mouse) mapping to 1 E4.

SOURCE

GP-39 (S-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of GP-39 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30465 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

GP-39 (S-18) is recommended for detection of GP-39 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


GP-39 (S-18) is also recommended for detection of GP-39 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for GP-39 siRNA (h): sc-44580, GP-39 siRNA (m): sc-44581, GP-39 shRNA Plasmid (h): sc-44580-SH, GP-39 shRNA Plasmid (m): sc-44581-SH, GP-39 shRNA (h) Lentiviral Particles: sc-44580-V and GP-39 shRNA (m) Lentiviral Particles: sc-44581-V.


Molecular Weight of GP-39: 39 kDa.

Positive Controls: GP-39 (h): 293 Lysate: sc-110933.

DATA



GP-39 (S-18): sc-30465. Western blot analysis of GP-39 expression in non-transfected: sc-110760 (A) and human GP-39 transfected: sc-110933 (B) 293 whole cell lysates.



GP-39 (S-18): sc-30465. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic staining of hepatocytes.

SELECT PRODUCT CITATIONS

1. Di Rosa, M., et al. 2014. Determination of chitinases family during osteoclastogenesis. *Bone* 61C: 55-63.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



MONOS
Satisfaction
Guaranteed

Try **GP-39 (D-11): sc-393494** or **GP-39 (E-11): sc-376910**, our highly recommended monoclonal alternatives to GP-39 (S-18).