

CD40 (P-14): sc-30684

BACKGROUND

Resting B cells can be activated and clonally expanded into antibody-producing cells in response to a combination of cell contact and soluble signals provided by primed helper T (Th) cells. While cytokines IL-4 and IL-13 alone are inadequate for B cell activation, contact with Th cells seems to be sufficient for delivery of proliferative signals. A receptor ligand pair central to the transmission of this signal is CD40, expressed on the surface of B cells, together with CD40L, expressed on activated T cells. In the presence of such stimulus, IL-4 and IL-13 are capable of triggering immunoglobulin class switching and secretion of IgE. B cells are sensitive to these cytokines only subsequent to CD40/CD40L-driven DNA synthesis. A downstream mediator of the CD40 signaling pathway, designated CRAF, is a member of an expanding family of proteins that contain a conserved cysteine- and histidine-rich RING finger motif. Other members of the family include TRAF1 and TRAF2. The latter proteins have been shown to regulate TNF-R2 as well as CD40 signaling through activation of the NF κ B family of transcription factors.

REFERENCES

1. Kehry, M.R., et al. 1994. B-cell activation by helper T-cell membranes. *Crit. Rev. Immunol.* 14: 221-238.
2. Hu, H.M., et al. 1994. A novel RING finger protein interacts with the cytoplasmic domain of CD40. *J. Biol. Chem.* 269: 30069-30072.
3. Rothe, M., et al. 1994. A novel family of putative signal transducers associated with the cytoplasmic domain of the 75 kDa tumor necrosis factor receptor. *Cell* 78: 681-682.
4. Gordon, J. 1995. CD40 and its ligand: central players in B lymphocyte survival, growth, and differentiation. *Blood Rev.* 9: 53-56.
5. Fuleihan, R., et al. 1995. Expression of the CD40 ligand in T lymphocytes and induction of IgE isotype switching. *Intl. Arch. All. Immunol.* 107: 43-44.

CHROMOSOMAL LOCATION

Genetic locus: Cd40 (mouse) mapping to 2 H3.

SOURCE

CD40 (P-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal extracellular domain of CD40 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30684 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CD40 (P-14) is recommended for detection of precursor and mature CD40 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CD40 siRNA (m): sc-29998, CD40 shRNA Plasmid (m): sc-29998-SH and CD40 shRNA (m) Lentiviral Particles: sc-29998-V.

Molecular Weight of CD40: 43 kDa.

Positive Controls: WEHI-231 whole cell lysate: sc-2213.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **CD40 (HM40-3): sc-20010**, our highly recommended monoclonal alternative to CD40 (P-14). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **CD40 (HM40-3): sc-20010**.