

SH-PTP2 (G-15): sc-30689

BACKGROUND

The steady state of protein tyrosyl phosphorylation in cells is regulated by the opposing action of tyrosine kinases and protein tyrosine phosphatases (PTPs). Several groups have independently identified a non-transmembrane PTP, designated SH-PTP1 (also known as PTP1C, HCP and SHP), which is primarily expressed in hematopoietic cells and characterized by the presence of two SH2 domains N-terminal to the PTP domain. SH2 domains generally mediate the association of regulatory molecules with specific phosphotyrosine-containing sites on autophosphorylated receptors, thereby controlling the initial interaction of receptors with these substrates. A second and much more widely expressed PTP with SH2 domains, SH-PTP2 (also designated PTP1D and Syp), has been identified. Strong sequence similarity between SH-PTP2 and the *Drosophila* gene corkscrew (CSW) and their similar patterns of expression suggest that SH-PTP2 is the human corkscrew homolog.

REFERENCES

1. Chernoff, J., et al. 1990. Cloning of a cDNA for a major human protein-tyrosine-phosphatase. *Proc. Natl. Acad. Sci. USA* 87: 2735-2739.
2. Shen, S., et al. 1991. A protein-tyrosine phosphatase with sequence similarity to the SH2 domain of the protein-tyrosine kinases. *Nature* 352: 736-739.
3. Plutzky, J., et al. 1992. Isolation of a Src homology 2-containing tyrosine phosphatase. *Proc. Natl. Acad. Sci. USA* 89: 1123-1127.
4. Yi, T., et al. 1992. Protein tyrosine phosphatase containing SH2 domains: characterization, preferential expression in hematopoietic cells, and localization to human chromosome 12p12-p13. *Mol. Cell. Biol.* 12: 836-846.
5. Matthews, R.J., et al. 1992. Characterization of hematopoietic intracellular protein tyrosine phosphatases: description of a phosphatase containing an SH2 domain and another enriched in proline-, glutamic acid-, serine-, and threonine-rich sequences. *Mol. Cell. Biol.* 12: 2396-2405.
6. Freeman, R.M., Jr., et al. 1992. Identification of a human src homology 2-containing protein-tyrosine-phosphatase: a putative homolog of *Drosophila* corkscrew. *Proc. Natl. Acad. Sci. USA* 89: 11239-11243.

CHROMOSOMAL LOCATION

Genetic locus: PTPN11 (human) mapping to 12q24.13; Ptpn11 (mouse) mapping to 5 F.

SOURCE

SH-PTP2 (G-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of SH-PTP2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30689 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SH-PTP2 (G-15) is recommended for detection of SH-PTP2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


SH-PTP2 (G-15) is also recommended for detection of SH-PTP2 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for SH-PTP2 siRNA (h): sc-36488, SH-PTP2 siRNA (m): sc-36489, SH-PTP2 shRNA Plasmid (h): sc-36488-SH, SH-PTP2 shRNA Plasmid (m): sc-36489-SH, SH-PTP2 shRNA (h) Lentiviral Particles: sc-36488-V and SH-PTP2 shRNA (m) Lentiviral Particles: sc-36489-V.


Molecular Weight of SH-PTP2: 70 kDa.

Positive Controls: SH-PTP2 (m2): 293T Lysate: sc-123530, Jurkat whole cell lysate: sc-2204 or NIH/3T3 whole cell lysate: sc-2210.

DATA



SH-PTP2 (G-15): sc-30689. Western blot analysis of SH-PTP2 expression in non-transfected: sc-117752 (A) and mouse SH-PTP2 transfected: sc-123530 (B) 293T whole cell lysates.



SH-PTP2 (G-15): sc-30689. Immunofluorescence staining of methanol-fixed NIH/3T3 cells showing cytoplasmic localization.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **SH-PTP2 (B-1): sc-7384** or **SH-PTP2 (D-3): sc-271053**, our highly recommended monoclonal alternatives to SH-PTP2 (G-15). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **SH-PTP2 (B-1): sc-7384**.