

Annexin VI (E-20): sc-30763

BACKGROUND

The Annexin family of calcium-binding proteins is composed of at least ten mammalian genes. It is characterized by a conserved core domain, which binds to phospholipids in a Ca^{2+} -dependent manner, and a unique amino terminal region, which may confer binding specificity. The Annexin family has been implicated as regulators of such diverse processes as ion-flux, endocytosis and exocytosis, and cellular adhesion. When overexpressed in A431 cells, Annexin VI causes a partial reversal of the transformed phenotype. It has been hypothesized that growth-dependent post-translational modifications of Annexins are required for proper sub-cellular localization. Annexin VII, also referred to as synexin, is located at the plasma membrane in normal muscle tissue. However, in muscle samples from patients suffering from Duchenne's muscular dystrophy, Annexin VII, along with Annexins IV and VI, are released into the cytoplasm and later, as the disease progresses, into the extracellular space. Two forms of Annexin XI, designated A and B, have been identified. Transfection of COS-7 cells with Annexin XI-A, but not Annexin XI-B, causes formation of Annexin XI-associated vesicles.

CHROMOSOMAL LOCATION

Genetic locus: ANXA6 (human) mapping to 5q33.1; Anxa6 (mouse) mapping to 11 B1.3.

SOURCE

Annexin VI (E-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Annexin VI of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30763 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Annexin VI (E-20) is recommended for detection of Annexin VI of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Annexin VI (E-20) is also recommended for detection of Annexin VI in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Annexin VI siRNA (h): sc-29688, Annexin VI siRNA (m): sc-29689, Annexin VI shRNA Plasmid (h): sc-29688-SH, Annexin VI shRNA Plasmid (m): sc-29689-SH, Annexin VI shRNA (h) Lentiviral Particles: sc-29688-V and Annexin VI shRNA (m) Lentiviral Particles: sc-29689-V.


Molecular Weight of Annexin VI: 68 kDa.

Positive Controls: L8 cell lysate: sc-3807, JAR cell lysate: sc-2276 or Annexin VI (h): 293 Lysate: sc-113205.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Annexin VI (E-20): sc-30763. Western blot analysis of Annexin VI expression in non-transfected: sc-110760 (A) and human Annexin VI transfected: sc-113205 (B) 293 whole cell lysates.

SELECT PRODUCT CITATIONS

1. Sui, J., et al. 2008. Comparative proteomics analysis of vascular smooth muscle cells incubated with S- and R-enantiomers of atenolol using iTRAQ-coupled two-dimensional LC-MS/MS. Mol. Cell. Proteomics 7: 1007-1018.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Annexin VI (E-5): sc-271859** or **Annexin VI (G-10): sc-166807**, our highly recommended monoclonal alternatives to Annexin VI (E-20).