

Cortactin (F-14): sc-30770

BACKGROUND

Cortactin (also designated Ems-1) is a filamentous actin (F-actin) binding protein that is present in 80 and 85 kDa forms. It has also been shown to be a substrate for Src p60. Cortactin contains tandem 37 amino acid repeats at the amino terminus and an SH3 domain at the carboxy terminus. The tandem repeats appear to be necessary for F-actin binding. Tyrosine phosphorylation of cortactin by Src p60 results in diminished F-actin binding to cortactin and reduced F-actin cross-linking activity. Cortactin has also been shown to be phosphorylated in response to FGF-1. Cortactin exhibits abundant expression in megakaryocytes and platelets, and it may play a role in the maturation of megakaryocytes.

REFERENCES

1. Wu, H., et al. 1993. Cortactin, an 80/85-kilodalton pp60src substrate, is a filamentous actin-binding protein enriched in the cell cortex. *J. Cell Biol.* 120: 1417-1426.
2. Zhan, X., et al. 1993. Murine cortactin is phosphorylated in response to fibroblast growth factor-1 on tyrosine residues late in the G₁ phase of the BALB/c 3T3 cell cycle. *J. Biol. Chem.* 268: 24427-24431.

CHROMOSOMAL LOCATION

Genetic locus: CTTN (human) mapping to 11q13; Ctnn (mouse) mapping to 7 F5.

SOURCE

Cortactin (F-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Cortactin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30770 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Cortactin (F-14) is recommended for detection of Cortactin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Cortactin siRNA (h): sc-35093, Cortactin siRNA (m): sc-35094, Cortactin shRNA Plasmid (h): sc-35093-SH, Cortactin shRNA Plasmid (m): sc-35094-SH, Cortactin shRNA (h) Lentiviral Particles: sc-35093-V and Cortactin shRNA (m) Lentiviral Particles: sc-35094-V.


Molecular Weight of Cortactin: 80 kDa.

Positive Controls: PC-12 cell lysate: sc-2250, A-431 whole cell lysate: sc-2201 or Cortactin (m): 293T Lysate: sc-126659.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Cortactin (F-14): sc-30770. Western blot analysis of Cortactin expression in non-transfected: sc-117752 (A) and mouse Cortactin transfected: sc-126659 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Cortactin (H-5): sc-55579** or **Cortactin (A-4): sc-55578**, our highly recommended monoclonal alternatives to Cortactin (F-14). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Cortactin (H-5): sc-55579**.