

UFD2 (N-16): sc-30838

BACKGROUND

UFD2, also designated ubiquitin conjugation factor E4 (UBE4B), binds to the ubiquitin moieties of preformed conjugates and catalyzes ubiquitin chain assembly in conjunction with E1, E2 and E3. During apoptosis, UFD2 is proteolytically cleaved at Asp 123 by caspase-6 and granzyme B, and is cleaved with approximately 10-fold less efficiency at Asp 109 by caspase-3 and caspase-7. In yeast, E4 activity is linked to cell survival under stress conditions, indicating that eukaryotes use E4-dependent proteolysis pathways for multiple cellular functions. In mammals, highest expression of UFD2 is in ovary, testis, heart and skeletal muscle.

REFERENCES

1. Koegl, M., et al. 1999. A novel ubiquitination factor, E4, is involved in multiubiquitin chain assembly. *Cell* 96: 635-644.
2. Conforti, L., et al. 2000. A Ufd2/D4Cole1e chimeric protein and overexpression of Rbp7 in the slow Wallerian degeneration (Wlds) mouse. *Proc. Natl. Acad. Sci. USA* 97: 11377-11382.
3. Krona, C., et al. 2003. Screening for gene mutations in a 500 kb neuroblastoma tumor suppressor candidate region in chromosome 1p; mutation and stage-specific expression in UBE4B/UFD2. *Oncogene* 22: 2343-2351.
4. Spinette, S., et al. 2004. Ufd2, a novel autoantigen in scleroderma, regulates sister chromatid separation. *Cell Cycle* 3: 1638-1644.
5. Saeki, Y., et al. 2004. Definitive evidence for Ufd2-catalyzed elongation of the ubiquitin chain through Lys48 linkage. *Biochem. Biophys. Res. Commun.* 320: 840-845.
6. Bazirgan, O.A., et al. 2005. Cdc48-Ufd2-Rad23: the road less ubiquitinated? *Nat. Cell Biol.* 7: 207-209.

CHROMOSOMAL LOCATION

Genetic locus: UBE4B (human) mapping to 1p36.22; Ube4b (mouse) mapping to 4 E2.

SOURCE

UFD2 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of UFD2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30838 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

UFD2 (N-16) is recommended for detection of UFD2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with UFD2/D4COLE1E (accession #10442019) fusion protein of mouse and rat origin.

UFD2 (N-16) is also recommended for detection of UFD2 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for UFD2 siRNA (h): sc-45980, UFD2 siRNA (m): sc-45981, UFD2 shRNA Plasmid (h): sc-45980-SH, UFD2 shRNA Plasmid (m): sc-45981-SH, UFD2 shRNA (h) Lentiviral Particles: sc-45980-V and UFD2 shRNA (m) Lentiviral Particles: sc-45981-V.


Molecular Weight of UFD2: 146 kDa.

Positive Controls: HeLa nuclear extract: sc-2120 or HS 181.Tes whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



UFD2 (N-16): sc-30838. Western blot analysis of UFD2 expression in HS 181.Tes whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.