

NF-YA (G-15): sc-30907

BACKGROUND

The CCAAT-binding factor NF-Y is a heteromeric transcription factor that specifically binds to CCAAT sequences in many eukaryotic genes. NF-Y is made up of three subunits, NF-YA, NF-YB and NF-YC. All three components are necessary for DNA binding. In each NF-Y subunit, the segment needed for formation of the NF-Y-DNA complex is conserved from yeast to human. These conserved segments are homologous to the histone-fold motif of eukaryotic histones. The DNA binding domains of the NF-YB and NF-YC subunits have been suggested to interact through a protein-protein histone-fold "handshake" motif in a manner analogous to the histone proteins, H2B and H2A, respectively.

REFERENCES

1. Baxeavanis, A.D., et al. 1995. A variety of DNA-binding and multimeric proteins contain the histone fold motif. *Nucleic Acids Res.* 23: 2685-2691.
2. Sinha, S., et al. 1996. Three classes of mutations in the A subunit of the CCAAT-binding factor CBF delineate functional domains involved in the three-step assembly of the CBF-DNA complex. *Mol. Cell. Biol.* 16: 328-337.
3. Currie, R.A. 1997. Functional interaction between the DNA binding subunit trimerization domain of NF-Y and high mobility group protein HMG-I(Y). *J. Biol. Chem.* 272: 30880-30888.
4. Maity, S.N., et al. 1998. Role of the CCAAT-binding protein CBF/NF-Y in transcription. *Trends Biochem. Sci.* 23: 174-178.
5. Liang, S.G., et al. 1998. Pathway of complex formation between DNA and three subunits of CBF/NF-Y. Photocross-linking analysis of DNA-protein interaction and characterization of equilibrium steps of subunit interaction and DNA binding. *J. Biol. Chem.* 273: 31590-31598.

CHROMOSOMAL LOCATION

Genetic locus: NFYA (human) mapping to 6p21.1; Nfya (mouse) mapping to 17 C.

SOURCE

NF-YA (G-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of NF-YA of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30907 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

NF-YA (G-15) is recommended for detection of NF-YA of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NF-YA (G-15) is also recommended for detection of NF-YA in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for NF-YA siRNA (h): sc-29947, NF-YA siRNA (m): sc-29948, NF-YA shRNA Plasmid (h): sc-29947-SH, NF-YA shRNA Plasmid (m): sc-29948-SH, NF-YA shRNA (h) Lentiviral Particles: sc-29947-V and NF-YA shRNA (m) Lentiviral Particles: sc-29948-V.


Molecular Weight of NF-YA: 30-50 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, NIH/3T3 whole cell lysate: sc-2210 or A-431 whole cell lysate: sc-2201.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



NF-YA (G-15): sc-30907. Western blot analysis of NF-YA expression in K-562 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **NF-YA (G-2): sc-17753**, our highly recommended monoclonal alternative to NF-YA (G-15). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **NF-YA (G-2): sc-17753**.