

mtTFA (A-16): sc-30962

BACKGROUND

mtTFA (mitochondrial transcription factor A), also known as mtTF1, TFAM, TCF6 (for transcription factor 6-like1), TCF6L2 and tsHMG, is a nuclear-encoded gene product that is imported into the mitochondria. mtTFA is required for many aspects of mitochondrial biogenesis including the replication and transcription of mitochondrial DNA (mtDNA). In mouse, testis-specific mtTFA is missing the mitochondria targeting sequence and is present in the nucleus rather than the mitochondria. This form of mtTFA is located primarily in the nuclei of elongated spermatids and may be involved in the regulation of gene expression of the haploid male genome. During mouse and human spermatogenesis there is a reduction of mtTFA protein levels and a reduction in mtDNA copy number. These features may provide one of the mechanisms by which paternal mtDNA transmission is prevented. mtTFA has been associated with mitochondrial disorder in humans characterized by ocular myopathy, exercise intolerance and muscle wasting.

REFERENCES

- Boissonneault, G., et al. 1993. A testis-specific gene encoding a nuclear high-mobility-group box protein located in elongating spermatids. *Mol. Cell. Biol.* 13: 4323-4330.
- Tiranti, V., et al. 1995. Chromosomal localization of mitochondrial transcription factor A (TCF6), single-stranded DNA-binding protein (SSBP), and endonuclease G (ENDOG), three human housekeeping genes involved in mitochondrial biogenesis. *Genomics* 25: 559-564.
- Larsson, N., et al. 1996. A single mouse gene encodes the mitochondrial transcription factor A and a testis-specific nuclear HMG-box protein. *Nat. Genet.* 13: 296-302.

CHROMOSOMAL LOCATION

Genetic locus: TFAM (human) mapping to 10q21.1; Tfam (mouse) mapping to 10 B5.3.

SOURCE

mtTFA (A-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of mtTFA of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30962 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS


mtTFA (A-16) is recommended for detection of precursor and mature mtTFA of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for mtTFA siRNA (h): sc-38053, mtTFA siRNA (m): sc-45912, mtTFA shRNA Plasmid (h): sc-38053-SH, mtTFA shRNA Plasmid (m): sc-45912-SH, mtTFA shRNA (h) Lentiviral Particles: sc-38053-V and mtTFA shRNA (m) Lentiviral Particles: sc-45912-V.


Molecular Weight of mtTFA: 25 kDa.

Positive Controls: CCRF-CEM cell lysate: sc-2225, A-431 whole cell lysate: sc-2201 or HeLa whole cell lysate: sc-2200.

DATA



mtTFA (A-16): sc-30962. Western blot analysis of mtTFA expression in A-431 whole cell lysate.



mtTFA (A-16): sc-30962. Immunofluorescence staining of methanol-fixed A-431 cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human esophagus tissue showing cytoplasmic staining of squamous epithelial cells (B).

SELECT PRODUCT CITATIONS

- Sastre-Serra, J., et al. 2012. The effects of 17β-estradiol on mitochondrial biogenesis and function in breast cancer cell lines are dependent on the ERα/ERβ ratio. *Cell. Physiol. Biochem.* 29: 261-268.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
Satisfation
Guaranteed

Try **mtTFA (C-9): sc-376672** or **mtTFA (F-6): sc-166965**, our highly recommended monoclonal alternatives to mtTFA (A-16). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **mtTFA (C-9): sc-376672**.