

Integrin α 8 (S-16): sc-30985

BACKGROUND

Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis. Integrin α 8 is expressed in contractile interstitial cells and smooth muscle cells and is upregulated in lung injury. Integrin α 8 is also a marker for smooth muscle cells, expressed as early as α smooth muscle actin. The Integrin α 8 chain is expressed in the glomerulus exclusively by mesangial cells and may play an important role for maintaining tissue integrity in the glomerulus during glomerular injury.

CHROMOSOMAL LOCATION

Genetic locus: ITGA8 (human) mapping to 10p13; Itga8 (mouse) mapping to 2 A1.

SOURCE

Integrin α 8 (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal extracellular domain of Integrin α 8 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-30985 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Integrin α 8 (S-16) is recommended for detection of Integrin α 8 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Integrin α 8 (S-16) is also recommended for detection of Integrin α 8 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Integrin α 8 siRNA (h): sc-35688, Integrin α 8 siRNA (m): sc-35689, Integrin α 8 shRNA Plasmid (h): sc-35688-SH, Integrin α 8 shRNA Plasmid (m): sc-35689-SH, Integrin α 8 shRNA (h) Lentiviral Particles: sc-35688-V and Integrin α 8 shRNA (m) Lentiviral Particles: sc-35689-V.

Molecular Weight of nonreduced Integrin α 8: 180 kDa.


Molecular Weight of reduced Integrin α 8: 155/25 kDa.

Positive Controls: Ramos cell lysate: sc-2216 or NAMALWA cell lysate: sc-2234.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



Integrin α 8 (S-16): sc-30985. Immunoperoxidase staining of formalin fixed, paraffin-embedded human kidney tissue showing membrane and cytoplasmic staining of cells in glomeruli and cytoplasmic staining of cells in tubules.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Integrin α 8 (F-11): sc-365798**, our highly recommended monoclonal alternative to Integrin α 8 (S-16).