

# MMP-3 (Y-16): sc-31074

## BACKGROUND

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including Collagen, gelatin, Fibronectin, Laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-3, MMP-10 and MMP-11 (also designated stromelysin-1, 2 and 3, respectively) activate procollagenase. MMP-3 activation of procollagenase can occur via two pathways. Direct activation by MMP-3 is slow and activation by MMP-3 in conjunction with tissue or plasma proteinases is rapid. MMP-10 is expressed in small intestine, and at lower levels in lung and heart. MMP-11 is specifically expressed in stromal cells of breast carcinomas and contributes to epithelial cell malignancies.

## REFERENCES

1. Saus, J., et al. 1988. The complete primary structure of human matrix metalloproteinase-3. Identity with stromelysin. *J. Biol. Chem.* 263: 6742-6745.
2. Basset, P., et al. 1990. A novel metalloproteinase gene specifically expressed in stromal cells of breast carcinomas. *Nature* 348: 699-704.
3. Suzuki, K., et al. 1990. Mechanisms of activation of tissue procollagenase by matrix metalloproteinase 3 (stromelysin). *Biochemistry* 29: 10261-10270.
4. Birkedal-Hansen, et al. 1993. Matrix metalloproteinases: a review. *Crit. Rev. Oral. Biol. Med.* 4: 197-250.
5. Reinemer, P., et al. 1994. Structural implications for the role of the N-terminus in the 'superactivation' of collagenases. A crystallographic study. *FEBS Lett.* 338: 227-233.
6. Knäuper, V., et al. 1996. Activation of human neutrophil procollagenase by stromelysin 2. *Eur. J. Biochem.* 235: 187-191.
7. Machein, U., et al. 1997. Expression of several matrix metalloproteinase genes in human monocytic cells. *Adv. Exp. Med. Biol.* 421: 247-251.
8. Madlener, M., et al. 1997. cDNA cloning and expression of the gene encoding murine stromelysin-2 (MMP-10). *Gene* 202: 75-81.

## CHROMOSOMAL LOCATION

Genetic locus: *Mmp3* (mouse) mapping to 9 A1.

## SOURCE

MMP-3 (Y-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MMP-3 of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31074 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

MMP-3 (Y-16) is recommended for detection of MMP-3 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MMP-3 siRNA (m): sc-37265, MMP-3 siRNA (r): sc-61874, MMP-3 shRNA Plasmid (m): sc-37265-SH, MMP-3 shRNA Plasmid (r): sc-61874-SH, MMP-3 shRNA (m) Lentiviral Particles: sc-37265-V and MMP-3 shRNA (r) Lentiviral Particles: sc-61874-V.


Molecular Weight of MMP-3: 55 kDa.

Positive Controls: rat placenta extract: sc-364808.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



MMP-3 (Y-16): sc-31074. Western blot analysis of mouse recombinant MMP-3.



MMP-3 (Y-16): sc-31074. Western blot analysis of MMP-3 expression in rat placenta tissue extract.

## SELECT PRODUCT CITATIONS

1. Kavas, A., et al. 2013. Potential of raloxifene in reversing osteoarthritis-like alterations in rat chondrocytes: an *in vitro* model study. *J. Biosci.* 38: 135-147.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.