

cyclin A (T-14): sc-31084

BACKGROUND

The critical role that the family of regulatory proteins known as cyclins play in eukaryotic cell cycle regulation is well established. The best-characterized cyclin complex is the mitotic cyclin B/Cdc2 p34 kinase, the active component of maturing promoting factor. Cyclin A accumulates prior to cyclin B in the cell cycle, appears to be involved in control of S phase and has been shown to associate with cyclin-dependent kinase-2 (Cdk2). In addition, cyclin A has been implicated in cell transformation and is found in complexes with E1A, transcription factors DRTF1 and E2F and retinoblastoma protein, p110. A second form of cyclin A, named cyclin A1 because of its high sequence homology to *Xenopus* cyclin A1, is most highly expressed in germ cells. It has been proposed that cyclin A1 can associate with Cdk2, p39 and Cdc2 p34.

REFERENCES

1. Draetta, G., et al. 1989. Cdc2 protein kinase is complexed with both cyclin A and B: evidence for proteolytic inactivation of MPF. *Cell* 56: 829-838.
2. Giordano, A., et al. 1989. A 60 kDa Cdc2-associated polypeptide complexes with the E1A proteins in adenovirus-infected cells. *Cell* 58: 981-990.
3. Gautier, J., et al. 1990. Cyclin is a component of maturation-promoting factor from *Xenopus*. *Cell* 60: 487-494.
4. Wang, J., et al. 1990. Hepatitis B virus integration in a cyclin A gene in a hepatocellular carcinoma. *Nature* 343: 555-557.
5. Pines, J., et al. 1990. Human cyclin A is adenovirus E1A-associated protein p60 and behaves differently from cyclin B. *Nature* 346: 760-763.
6. Bandara, L.R., et al. 1991. Cyclin A and the retinoblastoma gene product complex with a common transcription factor. *Nature* 352: 249-251.
7. Williams, R.T., et al. 1992. Co-purification of p34Cdc2/p58 cyclin A proline-directed protein kinase and the retinoblastoma tumor susceptibility gene product: interaction of an oncogenic Serine/Threonine protein kinase with a tumor-suppressor protein. *Oncogene* 7: 423-432.
8. Lees, E., et al. 1992. Cyclin E/Cdk2 and cyclin A/Cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev.* 6: 1874-1885.
9. Yang, R., et al. 1997. Characterization of a second human cyclin A that is highly expressed in testis and in several leukemic cell lines. *Cancer Res.* 57: 913-930.

CHROMOSOMAL LOCATION

Genetic locus: CCNA2 (human) mapping to 4q25-q31; Ccna2 (mouse) mapping to 3 B.

SOURCE

cyclin A (T-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of cyclin A of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31084 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

cyclin A (T-14) is recommended for detection of cyclin A of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 µg per 100–500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for cyclin A siRNA (h): sc-29282 and cyclin A siRNA (m): sc-29283.


Molecular Weight of cyclin A: 54 kDa.

Positive Controls: SK-BR-3 cell lysate: sc-2218, K-562 whole cell lysate: sc-2203 or MCF7 whole cell lysate: sc-2206.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



cyclin A (T-14): sc-31084. Western blot analysis of cyclin A expression in K-562 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.