

Hemoglobin γ (V-15): sc-31118

BACKGROUND

Hemoglobin (Hgb) is coupled to four iron-binding, methene-linked tetrapyrrole rings (heme). The α (16p13.3; 5'- ζ -pseudo ζ -pseudo α 2-pseudo α 1- α 2- α 1- θ 1-3') and β (11p15.5) globin loci determine the basic hemoglobin structure. The globin portion of hemoglobin consists of two α chains and two β chains arranged in pairs forming a tetramer. Each of the four globin chains covalently associates with a heme group. The bonds between α and β chains are weaker than between similar globin chains, thereby forming a cleavage plane that is important for oxygen binding and release. High affinity for oxygen occurs upon relaxation of the α 1- β 2 cleavage plane. When the two α 1- β 2 interfaces are closely bound, hemoglobin has a low affinity for oxygen. Hb A, which contains two α chains plus two β chains, comprises 97% of total circulating hemoglobin. The remaining 3% of total circulating hemoglobin is comprised of Hb A-2, which consists of two α chains plus two δ chains, and fetal hemoglobin (Hb F), which consists of two α chains together with two γ chains.

REFERENCES

- Liebhaber, S.A., Goossens, M. and Kan, Y.W. 1981. Homology and concerted evolution at the α 1 and α 2 loci of human α -globin. *Nature* 290: 26-29.
- Goodbourn, S.E., Higgs, D.R., Clegg, J.B. and Weatherall, D.J. 1983. Molecular basis of length polymorphism in the human ζ -globin gene complex. *Proc. Natl. Acad. Sci. USA* 80: 5022-5026.
- Giardina, B., Messina, I., Scatena, R. and Castagnola, M. 1995. The multiple functions of hemoglobin. *Crit. Rev. Biochem. Mol. Biol.* 30: 165-196.
- Adachi, K., Zhao, Y. and Surrey, S. 2002. Assembly of human hemoglobin (Hb) β - and γ -globin chains expressed in a cell-free system with α -globin chains to form Hb A and Hb F. *J. Biol. Chem.* 277: 13415-13420.
- Sudha, R., Anantharaman, L., Sivaram, M.V., Mirsamadi, N., Choudhury, D., Lohiya, N.K., Gupta, R.B. and Roy, R.P. 2004. Linkage of interactions in sickle hemoglobin fiber assembly: inhibitory effect emanating from mutations in the AB region of the α chain is annulled by a mutation at its EF corner. *J. Biol. Chem.* 279: 20018-20027.
- Feng, L., Gell, D.A., Zhou, S., Gu, L., Kong, Y., Li, J., Hu, M., Yan, N., Lee, C., Rich, A.M., Armstrong, R.S., Lay, P.A., Gow, A.J., Weiss, M.J., Mackay, J.P. and Shi, Y. 2004. Molecular mechanism of AHSP-mediated stabilization of α -hemoglobin. *Cell* 119: 629-640.
- Baudin-Creuzat, V., Vasseur-Godbillon, C., Pato, C., Prehu, C., Wajcman, H. and Marden, M.C. 2004. Transfer of human α - to β -hemoglobin via its chaperone protein: evidence for a new state. *J. Biol. Chem.* 279: 36530-36533.

CHROMOSOMAL LOCATION

Genetic locus: HBG1/HBG2/HBB/HBE1 (human) mapping to 11p15.4.

SOURCE

Hemoglobin γ (V-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Hemoglobin γ of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31118 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Hemoglobin γ (V-15) is recommended for detection of Hemoglobin γ and, to a lesser extent, hemoglobin β and hemoglobin ϵ of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Hemoglobin γ : 18 kDa.

Positive Controls: HEL 92.1.7 cell lysate: sc-2270 or TF-1 cell lysate: sc-2412.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Hemoglobin γ (51-7): sc-21756** or **Hemoglobin γ (B-4): sc-377395**, our highly recommended monoclonal alternatives to Hemoglobin γ (V-15).