PMS1 (I-16): sc-31143



The Power to Question

BACKGROUND

The finding that mutations in DNA mismatch repair genes are associated with hereditary nonpolyposis colorectal cancer (HNPCC) has resulted in considerable interest in the understanding of the mechanism of DNA mismatch repair. Initially, inherited mutations in the MSH2 and MLH1 homologs of the bacterial DNA mismatch repair genes MutS and MutL were demonstrated at high frequency in HNPCC and were shown to be associated with microsatellite instability. The demonstration that 10 to 45% of pancreatic, gastric, breast, ovarian and small cell lung cancers also display microsatellite instability has been interpreted to suggest that DNA mismatch repair is not restricted to HNPCC tumors but is a common feature in tumor initiation or progression. Two additional homologs of the prokaryotic MutL gene, designated PMS1 and PMS2, have been identified and shown to be mutated in the germline of HNPCC patients.

REFERENCES

- 1. Peltomäki, P., et al. 1993. Genetic mapping of a locus predisposing to human colorectal cancer. Science 260: 810-812.
- Ionov, Y., et al. 1993. Ubiquitous somatic mutations in simple repeated sequences reveal a new mechanism for colonic carcinogenesis. Nature 363: 558-561.
- 3. Papadopoulos, N., et al. 1994. Mutation of a MutL homolog in hereditary colon cancer. Science 263: 1625-1629.
- 4. Prolla, T.A., et al. 1994. MLH1, Pms1, and Msh2 interactions during the initation of DNA mismatch repair in yeast. Science 265: 1091-1092.
- 5. Palombo, F., et al. 1994. Mismatch repair and cancer. Nature 367: 417-418.
- Bronner, C.E., et al. 1994. Mutation in the DNA mismatch repair gene homologue hMLH1 is associated with hereditary non-polyposis colon cancer. Nature 368: 258-261.
- Nicolaides, N.C., et al. 1994. Mutations of two PMS homologues in hereditary nonpolyposis colon cancer. Nature 371: 75-80.

CHROMOSOMAL LOCATION

Genetic locus: PMS1 (human) mapping to 2q32.2; Pms1 (mouse) mapping to 1 C1.1.

SOURCE

PMS1 (I-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of PMS1 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31142 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

PMS1 (I-16) is recommended for detection of PMS1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PMS1 (I-16) is also recommended for detection of PMS1 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for PMS1 siRNA (h): sc-37409, PMS1 siRNA (m): sc-45504, PMS1 shRNA Plasmid (h): sc-37409-SH, PMS1 shRNA Plasmid (m): sc-45504-SH, PMS1 shRNA (h) Lentiviral Particles: sc-37409-V and PMS1 shRNA (m) Lentiviral Particles: sc-45504-V.

Molecular Weight of PMS1: 115 kDa.

Positive Controls: T24 cell lysate: sc-2292.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com