

p130 (K-17): sc-31338

BACKGROUND

The human retinoblastoma gene product (Rb) plays an important role in the negative regulation of cell proliferation. The Rb family includes p107 and p130, which form complexes with E2F proteins and share a high degree of structural homology in the adenovirus E1A binding domain (i.e. "pocket region"), which plays a primary role in the function of these proteins. The Rb family members undergo cell cycle dependent phosphorylation during mid-G₁ to S phase transition, which is dependent upon the activity of cyclin D/Cdk4. In contrast to pRb and p107, p130 is also phosphorylated during G₀ and the early G₁ phase of the cell cycle. p130 is specifically phosphorylated on serine and threonine residues in cells arrested in G₀ by serum deprivation or density arrest, and these residues are clustered within a short co-linear region unique to p130 defined as the Loop.

REFERENCES

1. Kovesdi, I., et al. 1986. Identification of a cellular transcription factor involved in E1A transactivation. *Cell* 45: 219-228.
2. Chellappan, S., et al. 1991. The E2F transcription factor is a cellular target for the Rb protein. *Cell* 65: 1053-1061.
3. Chittenden, T., et al. 1991. The T/E1A-binding domain of the retinoblastoma product can interact selectively with a sequence-specific DNA-binding protein. *Cell* 65: 1073-1082.
4. Bandara, L., et al. 1991. Cyclin A and the retinoblastoma gene product complex with a common transcription factor. *Nature* 352: 249-251.
5. Helin, K., et al. 1992. A cDNA encoding a pRb-binding protein with properties of the transcription factor E2F. *Cell* 70: 337-350.
6. Kaelin, W.G., Jr., et al. 1992. Expression cloning of a cDNA encoding a retinoblastoma-binding protein with E2F-like properties. *Cell* 70: 351-364.
7. Nevins, J.R. 1992. E2F: a link between the Rb tumor suppressor protein and viral oncoproteins. *Science* 258: 424-429.
8. Mayol, X., et al. 1993. Cloning of a new member of the retinoblastoma gene family (pRb2) which binds to the E1A transforming domain. *Oncogene* 8: 2561-2566.

CHROMOSOMAL LOCATION

Genetic locus: RBL2 (human) mapping to 16q12.2; Rbl2 (mouse) mapping to 8 C5.

SOURCE

p130 (K-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of p130 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31338 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p130 (K-17) is recommended for detection of p130 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

p130 (K-17) is also recommended for detection of p130 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for p130 siRNA (h): sc-29425, p130 siRNA (m): sc-29426, p130 shRNA Plasmid (h): sc-29425-SH, p130 shRNA Plasmid (m): sc-29426-SH, p130 shRNA (h) Lentiviral Particles: sc-29425-V and p130 shRNA (m) Lentiviral Particles: sc-29426-V.

Molecular Weight of p130: 130 kDa.

Positive Controls: Jurkat nuclear extract: sc-2132, Jurkat whole cell lysate: sc-2204 or C32 nuclear extract: sc-2136.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **p130 (A-10): sc-374521** or **p130 (DCS-215): sc-53641**, our highly recommended monoclonal alternatives to p130 (K-17).