

GABA_A Rα5 (S-20): sc-31417

BACKGROUND

GAD-65 and GAD-67, glutamate decarboxylases function to catalyze the production of GABA (γ-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl⁻ conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABA_A) and metabotropic (GABA_B) receptors as well as a third class of receptors called GABA_C. Both GABA_A and GABA_C are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABA_A receptor family include GABA_A Rα1-6, GABA_A Rβ1-3, GABA_A Rγ1-3, GABA_A Rδ, GABA_A Rε, GABA_A Rρ1 and GABA_A Rρ2. The GABA_B family is composed of GABA_B R1α and GABA_B R1β. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2, and -3). The GABA transporters function to terminate GABA action.

REFERENCES

1. Nelson, H., et al. 1990. Cloning of the human brain GABA transporter. FEBS Lett. 269: 181-184.
2. Cherubini, E., et al. 1991. GABA: an excitatory transmitter in early post-natal life. Trends Neurosci. 14: 515-519.
3. Borden, L.A., et al. 1992. Molecular heterogeneity of the γ-aminobutyric acid (GABA) transport system. Cloning of two novel high affinity GABA transporters from rat brain. J. Biol. Chem. 267: 21098-21104.
4. Dirx, R., Jr., et al. 1995. Targeting of the 67 kDa isoform of glutamic acid decarboxylase to intracellular organelles is mediated by its interaction with the NH₂-terminal region of the 65 kDa isoform of glutamic acid decarboxylase. J. Biol. Chem. 270: 2241-2246.
5. Lukasiewicz, P.D. 1996. GABA_C receptors in the vertebrate retina. Mol. Neurobiol. 12: 181-194.

CHROMOSOMAL LOCATION

Genetic locus: GABRA5 (human) mapping to 15q12; Gabra5 (mouse) mapping to 7 B4.

SOURCE

GABA_A Rα5 (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of GABA_A Rα5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31417 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GABA_A Rα5 (S-20) is recommended for detection of GABA_A Rα5 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GABA_A Rα5 siRNA (h): sc-42433, GABA_A Rα5 siRNA (m): sc-42434, GABA_A Rα5 shRNA Plasmid (h): sc-42433-SH, GABA_A Rα5 shRNA Plasmid (m): sc-42434-SH, GABA_A Rα5 shRNA (h) Lentiviral Particles: sc-42433-V and GABA_A Rα5 shRNA (m) Lentiviral Particles: sc-42434-V.


Molecular Weight of GABA_A Rα5: 55 kDa.

Positive Controls: GABA_A Rα5 (h): 293T Lysate: sc-111120 or human brain frontal cortex tissue extract.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GABA_A Rα5 (S-20): sc-31417. Western blot analysis of GABA_A Rα5 expression in rat hippocampus (A), rat brain (B), mouse brain (C) and human brain frontal cortex (D) tissue extracts.



GABA_A Rα5 (S-20): sc-31417. Western blot analysis of GABA_A Rα5 expression in human GABA_A Rα5 transfected: sc-111120 (A) and non-transfected: sc-110760 (B) 293 whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **GABA_A Rα5 (S-20)** or **GABA_A Rα1-6 (E-8): sc-376282**, our highly recommended monoclonal alternatives to GABA_A Rα5 (S-20). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **GABA_A Rα5 (S-20)**.