

# TRPV6 (E-16): sc-31446

## BACKGROUND

The transient receptor potential (TRP) protein family consists of a diverse group of cation channels functioning in a variety of homeostatic and regulatory pathways. Four subfamilies exist, based on channel domain homology, not activating stimuli: C type (canonical or classical), V type (vanilloid receptor related), M type (melastatin related) and P type (PKD). TRPV6 belongs to the V type subfamily, and it facilitates calcium entry across the plasma membrane in pancreas, placenta and to a lesser extent stomach and kidney tissue. Furthermore, prostate cancer cells overexpress TRPV6, while benign prostate tissues do not express the protein, implying a role for TRPV6 in malignant growth.

## REFERENCES

1. Nilius, B., et al. 2002. Fast and slow inactivation kinetics of the Ca<sup>2+</sup> channels ECaC1 and ECaC2 (TRPV5 and TRPV6). Role of the intracellular loop located between transmembrane segments 2 and 3. *J. Biol. Chem.* 277: 30852-30858.
2. Fixemer, T., et al. 2003. Expression of the Ca<sup>2+</sup>-selective cation channel TRPV6 in human prostate cancer: a novel prognostic marker for tumor progression. *Oncogene* 22: 7858-7861.
3. Birnbaumer, L., et al. 2003. A comparison of the genes coding for canonical TRP channels and their M, V and P relatives. *Cell Calcium* 33: 419-432.
4. Hirnet, D., et al. 2003. The TRPV6 gene, cDNA and protein. *Cell Calcium* 33: 509-518.
5. Hoenderop, J.G., et al. 2003. Homo- and heterotetrameric architecture of the epithelial Ca<sup>2+</sup> channels TRPV5 and TRPV6. *EMBO J.* 22: 776-785.
6. Wissenbach, U., et al. 2004. TRPV6 and prostate cancer: cancer growth beyond the prostate correlates with increased TRPV6 Ca<sup>2+</sup> channel expression. *Biochem. Biophys. Res. Commun.* 322: 1359-1363.
7. van de Graaf, S.F., et al. 2003. Functional expression of the epithelial Ca<sup>2+</sup> channels (TRPV5 and TRPV6) requires association of the S100A10-annexin 2 complex. *EMBO J.* 22: 1478-1487.
8. Lambers, T.T., et al. 2004. Regulation of the mouse epithelial Ca<sup>2+</sup> channel TRPV6 by the Ca<sup>2+</sup>-sensor calmodulin. *J. Biol. Chem.* 279: 28855-28861.

## CHROMOSOMAL LOCATION

Genetic locus: TRPV6 (human) mapping to 7q34; Trpv6 (mouse) mapping to 6 B2.1.

## SOURCE

TRPV6 (E-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of TRPV6 of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31445 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

TRPV6 (E-16) is recommended for detection of TRPV6 (also designated CaT1) of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TRPV6 (E-16) is also recommended for detection of TRPV6 (also designated CaT1) in additional species, including canine and porcine.

Suitable for use as control antibody for TRPV6 siRNA (h): sc-44171, TRPV6 siRNA (m): sc-44172, TRPV6 shRNA Plasmid (h): sc-44171-SH, TRPV6 shRNA Plasmid (m): sc-44172-SH, TRPV6 shRNA (h) Lentiviral Particles: sc-44171-V and TRPV6 shRNA (m) Lentiviral Particles: sc-44172-V.

Molecular Weight of TRPV6 core: 75 kDa

Molecular Weight of glycosylated TRPV6 85-100 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, Caco-2 cell lysate: sc-2262 or SW480 cell lysate: sc-2219.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.