

PLU-1 (N-19): sc-31612

BACKGROUND

PLU-1 is a large (1544 amino acids) nuclear protein that interacts with brain factor-1 (BF-1) and paired box 9 (PAX9), both of which are developmental transcription factors. PLU-1 belongs to the testis-cancer antigen group of proteins and is a member of the ARID family of DNA binding proteins. It is a multi-domain protein with strong transcriptional repression properties. PLU-1 shows restricted expression in adult tissues, with high expression in testis, and transiently in the pregnant mammary gland. Both the PLU-1 gene and the PLU-1 protein product are specifically up-regulated in breast cancer. PLU-1 may be important in meiotic transcription because of its apparent association with chromatin.

REFERENCES

- Lu, P.J., et al. 1999. A novel gene (PLU-1) containing highly conserved putative DNA/chromatin binding motifs is specifically up-regulated in breast cancer. *J. Biol. Chem.* 274: 15633-15645.
- Madsen, B., et al. 2002. Characterisation and developmental expression of mouse PLU-1, a homologue of a human nuclear protein (PLU-1) which is specifically up-regulated in breast cancer. *Mech. Dev.* 119S239-119S246.
- Tan, K., et al. 2003. Human PLU-1 Has transcriptional repression properties and interacts with the developmental transcription factors BF-1 and PAX9. *J. Biol. Chem.* 278: 20507-20513.
- Madsen, B., et al. 2003. PLU-1, a transcriptional repressor and putative testis-cancer antigen, has a specific expression and localisation pattern during meiosis. *Chromosoma.* 112: 124-132.
- Catteau, A., et al. 2004. A short region of the promoter of the breast cancer associated PLU-1 gene can regulate transcription *in vitro* and *in vivo*. *Int. J. Oncol.* 25: 5-16.

CHROMOSOMAL LOCATION

Genetic locus: PLU-1 (human) mapping to 1q32.1; Plu-1 (mouse) mapping to 1 E4.

SOURCE

PLU-1 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of PLU-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31612 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-31612 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

PLU-1 (N-19) is recommended for detection of PLU-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PLU-1 siRNA (h): sc-44522, PLU-1 siRNA (m): sc-44523, PLU-1 shRNA Plasmid (h): sc-44522-SH, PLU-1 shRNA Plasmid (m): sc-44523-SH, PLU-1 shRNA (h) Lentiviral Particles: sc-44522-V and PLU-1 shRNA (m) Lentiviral Particles: sc-44523-V.

PLU-1 (N-19) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **PLU-1 (JARIA3D4): sc-81358**, our highly recommended monoclonal alternative to PLU-1 (N-19).