

NTR2 (H-19): sc-31696

BACKGROUND

Neurotensin (NT) initiates an intracellular response by interacting with the G protein-coupled receptors NTR1 (NTS1 receptor, high affinity NTR) and NTR2 (NTS2 receptor, levocabastine-sensitive neurotensin receptor), and the type I receptor NTR3 (NTS3 receptor, sortilin-1, Gp95). NT has a wide distribution in regions of the brain and in peripheral tissues where NT receptors can contribute to hypotension, hyperglycemia, hypothermia, antinociception and regulation of intestinal motility and secretion. HL-60 cells express NTR1, which can couple to G_{q^*} , $G_{i/o}$ or G_s . Alternative splicing of rat NTR2 can generate a 5-transmembrane domain variant isoform that is coexpressed with the full length NTR2 throughout the brain and spinal cord. NTR3 activation in the murine microglial cell line N11 induces MIP-2, MCP-1, IL-1 β and TNF α in an ERK1/2 and Akt kinase-dependent manner.

REFERENCES

- Nielsen, M.S., et al. 1999. Sortilin/neurotensin receptor-3 binds and mediates degradation of lipoprotein lipase. *J. Biol. Chem.* 274: 8832-8836.
- Choi, S.Y., et al. 1999. Characterization of high affinity neurotensin receptor NTR1 in HL-60 cells and its downregulation during granulocytic differentiation. *Br. J. Pharmacol.* 126: 1050-1056.
- Navarro, V., et al. 2002. Shedding of the luminal domain of the neurotensin receptor-3/sortilin in the HT29 cell line. *Biochem. Biophys. Res. Commun.* 298: 760-764.
- Martin, S., et al. 2003. Involvement of the neurotensin receptor-3 in the neurotensin-induced migration of human microglia. *J. Neurosci.* 23: 1198-1205.
- Dicou, E., et al. 2004. Neurotensin receptor-3/sortilin mediates neurotensin-induced cytokine/chemokine expression in a murine microglial cell line. *J. Neurosci. Res.* 78: 92-99.
- Leonetti, M., et al. 2004. Specific involvement of neurotensin type 1 receptor in the neurotensin-mediated *in vivo* dopamine efflux using knock-out mice. *J. Neurochem.* 89: 1-6.
- Chen, L., et al. 2004. Neurotensin depolarizes globus pallidus neurons in rats via neurotensin type-1 receptor. *Neuroscience* 125: 853-859.
- Liu, F., et al. 2004. Neurotensin negatively modulates Akt activity in neurotensin receptor-1-transfected AV12 cells. *J. Cell. Biochem.* 92: 603-611.

CHROMOSOMAL LOCATION

Genetic locus: NTSR2 (human) mapping to 2p25.1; Ntsr2 (mouse) mapping to 12 A1.1.

SOURCE

NTR2 (H-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of NTR2 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31696 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NTR2 (H-19) is recommended for detection of neurotensin receptor type 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NTR2 (H-19) is also recommended for detection of neurotensin receptor type 2 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for NTR2 siRNA (h): sc-44483, NTR2 siRNA (m): sc-44484, NTR2 shRNA Plasmid (h): sc-44483-SH, NTR2 shRNA Plasmid (m): sc-44484-SH, NTR2 shRNA (h) Lentiviral Particles: sc-44483-V and NTR2 shRNA (m) Lentiviral Particles: sc-44484-V.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.