

# PFK-1 (T-18): sc-31710

## BACKGROUND

Phosphofructokinases (PFK) are regulatory glycolytic enzymes that convert fructose 6-phosphate and ATP into fructose 1,6-bisphosphate (through PFK-1), fructose 2,6-bisphosphate (through PFK-2), and ADP. Human PFK-1 is tetrameric and isoenzymes include, PFK-1 muscle (PFKM, PFK-A), PFK-1 liver (PFKL, PFK-B), and PFK-1 platelet (PFKP, PFK-C, PFKF). PFK-1 is inhibited by ATP and citrate (from the tricarboxylic acid cycle). PFK-1 undergoes activation in the presence of elevated AMP. The most potent activator is fructose-2,6-bisphosphate, which is produced by PFK-2 from the same substrate, fructose 6-phosphate. PFK-2 is bifunctional and a key regulator for PFK-1. PFK-2 catalyzes the synthesis of fructose-2,6-bisphosphate, and contains fructose-2,6-bisphosphatase activity that catalyzes the degradation of fructose-2,6-bisphosphate. PFK-2 is dimeric and isoenzymes include PFK-2 liver (PFKFB1, PFRX), PFK-2 cardiac (PFKFB2), PFK-2 placental (PFKFB3, inducible PFK-2) and PFK-2 testis (PFKFB4).

## REFERENCES

1. Tsuura, Y., et al. 1998. Endogenous nitric oxide inhibits glucose-induced Insulin secretion by suppression of phosphofructokinase activity in pancreatic islets. *Biochem. Biophys. Res. Commun.* 252: 34-38.
2. Chang, S.H., et al. 2002. Role of Ser 530, Arg 292, and His 662 in the allosteric behavior of rabbit muscle phosphofructokinase. *Biochem. Biophys. Res. Commun.* 290: 670-675.
3. Zeitschel, U., et al. 2002. Changes in activity and expression of phosphofructokinase in different rat brain regions after basal forebrain cholinergic lesion. *J. Neurochem.* 83: 371-380.
4. Su, Y., et al. 2003. The  $\alpha$  subunit of the V-type H<sup>+</sup>-ATPase interacts with phosphofructokinase-1 in humans. *J. Biol. Chem.* 278: 20013-20018.
5. Sotgia, F., et al. 2003. Phosphofructokinase muscle-specific isoform requires caveolin-3 expression for plasma membrane recruitment and caveolar targeting: implications for the pathogenesis of caveolin-related muscle diseases. *Am. J. Pathol.* 163: 2619-2634.

## CHROMOSOMAL LOCATION

Genetic locus: PFKM (human) mapping to 12q13.11; PfkM (mouse) mapping to 15 F1.

## SOURCE

PFK-1 (T-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of PFK-1 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31710 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

PFK-1 (T-18) is recommended for detection of muscle type PFK-1 of human, mouse and, to a lesser extent, rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with liver PFK-1.

PFK-1 (T-18) is also recommended for detection of muscle type PFK-1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PFK-1 siRNA (h): sc-44561, PFK-1 siRNA (m): sc-44562, PFK-1 shRNA Plasmid (h): sc-44561-SH, PFK-1 shRNA Plasmid (m): sc-44562-SH, PFK-1 shRNA (h) Lentiviral Particles: sc-44561-V and PFK-1 shRNA (m) Lentiviral Particles: sc-44562-V.


Molecular Weight of PFK-1: 85 kDa.

Positive Controls: rat skeletal muscle extract: sc-364810.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



PFK-1 (T-18): sc-31710. Western blot analysis of PFK-1 expression in rat skeletal muscle tissue extract.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

**MONOS**  
Satisfaction  
Guaranteed

Try **PFK-1 (E-4): sc-377346** or **PFK-1 (G-11): sc-166722**, our highly recommended monoclonal alternatives to PFK-1 (T-18).