

FOXP3 (K-13): sc-31739



The Power to Question

BACKGROUND

The FOX family of transcription factors is a large group of proteins that share a common DNA binding domain termed a winged-helix or forkhead domain. During early development, FOXP1 and FOXP2 are expressed abundantly in the lung, with lower levels of expression in neural, intestinal and cardiovascular tissues, where they act as transcription repressors. FOXP1 is widely expressed in adult tissues, while neoplastic cells often exhibit a dramatic change in expression level or localization of FOXP1. The gene encoding human FOXP1 maps to chromosome 3p14.1. The gene encoding human FOXP2 maps to chromosome 7q31. The gene encoding FOXP3, a third member of this family, maps to chromosome Xp11.23. Mutations in this gene cause IPEX, a fatal, X-linked inherited disorder characterized by immune dysregulation. The FOXP3 protein, also known as scurfin, is essential for normal immune homeostasis. Specifically, FOXP3 represses transcription through a DNA binding forkhead domain, thereby regulating T cell activation.

REFERENCES

- Lai, C.S., et al. 2000. The SPCH1 region on human 7q31: genomic characterization of the critical interval and localization of translocations associated with speech and language disorder. *Am. J. Hum. Genet.* 67: 357-368.
- Banham, A.H., et al. 2001. The FOXP1 winged helix transcription factor is a novel candidate tumor suppressor gene on chromosome 3p. *Cancer Res.* 61: 8820-8829.
- Bennett, C.L., et al. 2001. The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat. Genet.* 27: 20-21.

CHROMOSOMAL LOCATION

Genetic locus: FOXP3 (human) mapping to Xp11.23; Foxp3 (mouse) mapping to X A1.1.

SOURCE

FOXP3 (K-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of FOXP3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31739 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-31739 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

FOXP3 (K-13) is recommended for detection of FOXP3 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

FOXP3 (K-13) is also recommended for detection of FOXP3 in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for FOXP3 siRNA (h): sc-43569, FOXP3 siRNA (m): sc-45646, FOXP3 shRNA Plasmid (h): sc-43569-SH, FOXP3 shRNA Plasmid (m): sc-45646-SH, FOXP3 shRNA (h) Lentiviral Particles: sc-43569-V and FOXP3 shRNA (m) Lentiviral Particles: sc-45646-V.

FOXP3 (K-13) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of FOXP3: 48 kDa.

Positive Controls: mouse brain extract: sc-2253 or Jurkat whole cell lysate: sc-2204.

DATA



FOXP3 (K-13): sc-31739. Immunoperoxidase staining of formalin fixed, paraffin-embedded human tonsil tissue showing nuclear staining of cells in germinal center and cells in non-germinal center.

SELECT PRODUCT CITATIONS

- Xu, G., et al. 2009. Activated mammalian target of rapamycin is associated with T regulatory cell insufficiency in nasal polyps. *Respir. Res.* 10: 13.
- Ma, C., et al. 2011. Colorectal cancer-derived Foxp3⁺ IL-17⁺ T cells suppress tumour-specific CD8⁺ T cells. *Scand. J. Immunol.* 74: 47-51.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **FOXP3 (2A11G9): sc-53876** or **FOXP3 (F-9): sc-166212**, our highly recommended monoclonal alternatives to FOXP3 (K-13). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **FOXP3 (2A11G9): sc-53876**.