

Enolase (V-15): sc-31858

BACKGROUND

Enolases have been characterized as highly conserved cytoplasmic glycolytic enzymes that may be involved in differentiation. Three isoenzymes have been identified: α Enolase, β Enolase and γ Enolase. α Enolase expression has been detected on most tissues, whereas β Enolase is expressed predominantly in muscle tissue and γ Enolase is detected only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphoenolpyruvic acid in the glycolytic pathway. The 433 amino acid protein shows 67% homology to yeast Enolase and 94% homology to rat nonneural Enolase. Studies also indicate that α Enolase is encoded by the same gene that encodes τ -crystallin, a lens structural protein.

REFERENCES

- Whitehead, M.C., et al. 1982. Synapse formation is related to the onset of neuron-specific Enolase immunoreactivity in the avian auditory and vestibular systems. *Dev. Neurosci.* 5: 298-307.
- Giallongo, A., et al. 1986. Molecular cloning and nucleotide sequence of a full-length cDNA for human α Enolase. *Proc. Natl. Acad. Sci. USA* 83: 6741-6745.
- Wistow, G.J., et al. 1989. τ -crystallin/ α Enolase: one gene encodes both an enzyme and a lens structural protein. *J. Cell Biol.* 107: 2729-2736.
- Verma, M., et al. 1994. DNA sequences encoding Enolase are remarkably conserved from yeast to mammals. *Life Sci.* 55: 893-899.
- Keller, A., et al. 1994. Coexpression of α and γ Enolase genes in neurons of adult rat brain. *J. Neurosci. Res.* 38: 493-504.
- Zhang, E., et al. 1997. Mechanism of enolase: the crystal structure of asymmetric dimer enolase-2-phospho-D-glycerate/enolase-phosphoenolpyruvate at 2.0 Å resolution. *Biochemistry* 36: 12526-12534.
- Deloulme, J.C., et al. 1997. A comparative study of the distribution of α and γ Enolase subunits in cultured rat neural cells and fibroblasts. *Int. J. Dev. Neurosci.* 15: 183-194.

SOURCE

Enolase (V-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Enolase of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-31858 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Enolase (V-15) is recommended for detection of α Enolase, β Enolase and γ Enolase of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Enolase (V-15) is also recommended for detection of α Enolase, β Enolase and γ Enolase in additional species, including equine, canine, bovine, porcine and avian.


Molecular Weight of Enolase: 48 kDa.

Positive Controls: IMR-32 whole cell lysate: sc-2409, γ Enolase (h): 293T Lysate: sc-170262 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Enolase (V-15): sc-31858. Western blot analysis of Enolase expression in IMR-32 (A), Hep G2 (B), A549 (C), U-937 (D) and SH-SY5Y (E) whole cell lysates and mouse brain tissue extract (F).



Enolase (V-15): sc-31858. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Enolase (A-5): sc-271384** or **Enolase (D-8): sc-390163**, our highly recommended monoclonal alternatives to Enolase (V-15). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Enolase (A-5): sc-271384**.