

Ras GAP (D-18): sc-32047

BACKGROUND

The mammalian c-H-, c-K- and N-Ras proto-oncogenes encode ubiquitously expressed proteins. p21^{Ras} can exist in either a physiologically quiescent GDP-binding state or a GTP-binding signal-emitting state. Oncogenic p21^{Ras} proteins are trapped in the excited signal-emitting state because the mechanism normally employed to delimit their excitation period, hydrolysis of their bound GTP to GDP, is impaired as a result of specific mutations. Interaction of p21^{Ras} with GTPase activating protein (GAP) can increase hydrolysis of p21^{Ras}-bound GTP by as much as 1000-fold. The product of the neurofibromatosis type 1 gene (NF1) has also been shown to exhibit p21^{Ras} GAP activity, and proteins that stimulate the GTPase activity of three other low molecular weight GTPases, including Rho, Rab 3A and Rap 1, have also been described.

REFERENCES

- Shih, T.Y., et al. 1980. Guanine nucleotide- binding and autophosphorylating activities associated with the p21^{src} protein of Harvey murine sarcoma virus. *Nature* 287: 686-691.
- Barbacid, M. 1987. Ras genes. *Annu. Rev. Biochem.* 56: 779-827.
- Trahey, M., et al. 1988. Molecular cloning of two types of GAP complementary DNA from human placenta. *Science* 242: 1697-1700.
- Vogel, U.S., et al. 1988. Cloning of bovine GAP and its interaction with oncogenic Ras p21. *Nature* 335: 90-93.
- McCormick, F. 1989. ras GTPase activating protein: signal transmitter and signal terminator. *Cell* 56: 5-8.
- Ballester, R., et al. 1990. The NF1 locus encodes a protein functionally related to mammalian GAP and yeast IRA proteins. *Cell* 63: 851-859.
- Martin, G.A., et al. 1990. The GAP-related domain of the neurofibromatosis type 1 gene product interacts with Ras p21. *Cell* 63: 843-849.
- Rubinfeld, B., et al. 1991. Molecular cloning of a GTPase activating protein specific for the Krev-1 protein p21^{rap1}. *Cell* 65: 1033-1042.
- Diekmann, D., et al. 1991. Bcr encodes a GTPase-activating protein for p21^{rac}. *Nature* 351: 400-402.

CHROMOSOMAL LOCATION

Genetic locus: RASA1 (human) mapping to 5q14.3; Rasa1 (mouse) mapping to 13 C3.

SOURCE

Ras GAP (D-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Ras GAP of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32047 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Ras GAP (D-18) is recommended for detection of Ras GAP p120 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Ras GAP (D-18) is also recommended for detection of Ras GAP p120 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Ras GAP siRNA (h): sc-29467, Ras GAP siRNA (m): sc-36394, Ras GAP shRNA Plasmid (h): sc-29467-SH, Ras GAP shRNA Plasmid (m): sc-36394-SH, Ras GAP shRNA (h) Lentiviral Particles: sc-29467-V and Ras GAP shRNA (m) Lentiviral Particles: sc-36394-V.

Molecular Weight of Ras GAP: 120 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, KNRK whole cell lysate: sc-2214 or 3611-RF whole cell lysate: sc-2215.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE** Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Ras GAP (B4F8): sc-63** or **Ras GAP (D4B4): sc-64**, our highly recommended monoclonal alternatives to Ras GAP (D-18). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Ras GAP (B4F8): sc-63**.