

CKR-4 (N-19): sc-32132

BACKGROUND

C-C or β chemokine family members are characterized by a pair of adjacent cysteine residues and serve as potent chemoattractants and activators of monocytes and T cells. C-C chemokine receptor family members include CKR-1, CKR-2A, CKR-2B, CKR-3, CKR-4, CKR-5, CKR-6, CKR-7, CKR-8, CKR-9, CKR-10 and the Duffy blood group antigen. Each of these receptors are G protein-coupled, seven pass transmembrane domain proteins whose major physiological role is to function in the chemotaxis of T cells and phagocytic cells to areas of inflammation. However, this receptor family has also been shown to facilitate viral infection. CKR-4 (C-C chemokine receptor type 4), also known as CCR4 or CMKBR4, is a 360 amino acid multi-pass membrane protein that localizes to the cell membrane and belongs to the C-C chemokine receptor family. Expressed at high levels in peripheral blood leukocytes and thymus tissue, CKR-4 functions as a high affinity receptor for C-C type chemokines and is thought to be involved in hippocampal-neuron survival.

REFERENCES

- Schweickart, V.L., Raport, C.J., Godiska, R., Byers, M.G., Eddy, R.L., Jr., Shows, T.B. and Gray, P.W. 1994. Cloning of human and mouse EBI1, a lymphoid-specific G protein-coupled receptor encoded on human chromosome 17q12-q21.2. *Genomics* 23: 643-650.
- Deng, H., Liu, R., Ellmeier, W., Choe, S., Unutmaz, D., Burkhart, M., Di Marzio, P., Marmon, S., Sutton, R.E. and Hill, C.M. 1996. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 381: 661-666.
- Dragic, T., Litwin, V., Allaway, G.P., Martin, S.R., Huang, Y., Nagashima, K.A., Cayanan, C., Maddon, P.J., Koup, R.A. and Moore, J.P. 1996. HIV-1 entry into CD4⁺ cells is mediated by the chemokine receptor CC-CKR-5. *Nature* 381: 667-673.
- Feng, Y., Broder, C.C., Kennedy, P.E. and Berger, E.A. 1996. HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 272: 872-877.
- Alkhatib, G., Combadiere, C., Broder, C.C., Feng, Y., Kennedy, P.E., Murphy, P.M. and Berger, E.A. 1996. CC CKR5: a RANTES, MIP-1 α , MIP-1 β receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science* 272: 1955-1958.
- Choe, H., Farzan, M., Sun, Y., Sullivan, N., Rollins, B., Ponath, P.D., Wu, L., Mackay, C.R., LaRosa, G. and Newman, W. 1996. The β chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV-1 isolates. *Cell* 85: 1135-1148.
- Doranz, B.J., Rucker, J., Yi, Y., Smyth, R.J., Samson, M., Peiper, S.C., Parmentier, M., Collman, R.G. and Doms, R.W. 1996. A dual-tropic primary HIV-1 isolate that uses fusin and the β chemokine receptors CKR-5, CKR-3, and CKR-2B as fusion cofactors. *Cell* 85: 1149-1158.
- Baba, M., Imai, T., Nishimura, M., Kakizaki, M., Takagi, S., Hieshima, K., Nomiyama, H. and Yoshie, O. 1997. Identification of CCR6, the specific receptor for a novel lymphocyte-directed C-C chemokine LARC. *J. Biol. Chem.* 272: 14893-14898.

CHROMOSOMAL LOCATION

Genetic locus: CCR4 (human) mapping to 3p22.3.

SOURCE

CKR-4 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal extracellular domain of CKR-4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32132 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CKR-4 (N-19) is recommended for detection of CKR-4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CKR-4 (N-19) is also recommended for detection of CKR-4 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for CKR-4 siRNA (h): sc-39886, CKR-4 siRNA (m): sc-39887, CKR-4 shRNA Plasmid (h): sc-39886-SH, CKR-4 shRNA Plasmid (m): sc-39887-SH, CKR-4 shRNA (h) Lentiviral Particles: sc-39886-V and CKR-4 shRNA (m) Lentiviral Particles: sc-39887-V.

Molecular Weight of CKR-4: 41 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.