

HIF-3 α (T-15): sc-32142

BACKGROUND

Cell growth and viability is compromised by oxygen deprivation (hypoxia). Hypoxia-inducible factors, including HIF-1 α , HIF-1 β (also designated Arnt 1), EPAS-1 (also designated HIF-2 α) and HIF-3 α , induce glycolysis, erythropoiesis and angiogenesis in order to restore oxygen homeostasis. Hypoxia-inducible factors are members of the Per-Arnt-Sim (PAS) domain transcription factor family. In response to hypoxia, HIF-1 α is upregulated and forms a heterodimer with Arnt 1 to form the HIF-1 complex. The HIF-1 complex recognizes and binds to the hypoxia responsive element (HRE) of hypoxia-inducible genes, thereby activating transcription. Hypoxia-inducible expression of some genes such as Glut-1, p53, p21 or Bcl-2, is HIF-1 α dependent, whereas expression of others, such as p27, GADD 153 or HO-1, is HIF-1 α independent. EPAS-1 and HIF-3 α have also been shown to form heterodimeric complexes with Arnt 1 in response to hypoxia.

REFERENCES

1. Wang, G.L., Jiang, B.H., Rue, E.A. and Semenza, G.L. 1995. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc. Natl. Acad. Sci. USA* 92: 5510-5514.
2. Tian, H., McKnight, S.L. and Russell, D.W. 1997. Endothelial PAS domain protein 1 (EPAS1), a transcription factor selectively expressed in endothelial cells. *Genes Dev.* 11: 72-82.
3. Luo, G., Gu, Y.Z., Jain, S., Chan, W.K., Carr, K.M., Hogenesch, J.B. and Bradfield, C.A. 1997. Molecular characterization of the murine HIF-1 α locus. *Gene Expr.* 6: 287-299.
4. Carmeliet, P., Dor, Y., Herbert, J.M., Fukumura, D., Brusselmans, K., Dewerchin, M., Neeman, M., Bono, F., Abramovitch, R., Maxwell, P., Koch, C.J., Ratcliffe, P., Moons, L., Jain, R.K., Collen, D. and Keshet, E. 1998. Role of HIF-1 α in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature* 394: 485-490.
5. Gu, Y.Z., Moran, S.M., Hogenesch, J.B., Wartman, L. and Bradfield, C.A. 1998. Molecular characterization and chromosomal localization of a third α -class hypoxia inducible factor subunit, HIF-3 α . *Gene Expr.* 7: 205-213.

CHROMOSOMAL LOCATION

Genetic locus: HIF3A (human) mapping to 19q13.32; Hif3a (mouse) mapping to 7 A2.

SOURCE

HIF-3 α (T-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of HIF-3 α of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-32142 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-32142 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HIF-3 α (T-15) is recommended for detection of HIF-3 α of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HIF-3 α (T-15) is also recommended for detection of HIF-3 α in additional species, including canine, bovine and porcine.


Suitable for use as control antibody for HIF-3 α siRNA (h): sc-38167, HIF-3 α siRNA (m): sc-38168, HIF-3 α shRNA Plasmid (h): sc-38167-SH, HIF-3 α shRNA Plasmid (m): sc-38168-SH, HIF-3 α shRNA (h) Lentiviral Particles: sc-38167-V and HIF-3 α shRNA (m) Lentiviral Particles: sc-38168-V.

HIF-3 α (T-15) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of HIF-3 α : 73 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, mouse liver extract: sc-2256 or rat liver extract: sc-2395.

DATA



HIF-3 α (T-15): sc-32142. Western blot analysis of human recombinant HIF-3 α fusion protein.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **HIF-3 α (E-8): sc-390933** or **HIF-3 α (D-7): sc-390769**, our highly recommended monoclonal alternatives to HIF-3 α (T-15).