

CYP24 (C-18): sc-32166

BACKGROUND

P450 enzymes constitute a family of monooxygenase enzymes that are involved in the metabolism of a wide array of endogenous and xenobiotic compounds. P450 enzymes can be classified, based on their sequence similarities, into distinct subfamilies, which include CYP1A and CYP2A. The P450 family member CYP19 catalyzes the conversion of C19 steroids to estrogens in various tissues, including placenta, gonads, adipose tissue, skin and brain. P450 cholesterol 7 α -hydroxylase, CYP7A1, is the rate limiting enzyme of bile acid synthesis in the liver, and its expression is mediated by the bile acid receptor FXR. CYP27A1 catalyzes vitamin D 25-hydroxylation and is localized to the mitochondria in kidney and liver. Overexpression of CYP24 (encoding vitamin D 24-hydroxylase) is likely to lead to abrogation of growth control mediated by vitamin D.

CHROMOSOMAL LOCATION

Genetic locus: CYP24A1 (human) mapping to 20q13.2; Cyp24a1 (mouse) mapping to 2 H3.

SOURCE

CYP24 (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of CYP24 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32166 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CYP24 (C-18) is recommended for detection of mature CYP24 of human, mouse and, to a lesser extent, rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CYP24 (C-18) is also recommended for detection of mature CYP24 in additional species, including equine.

Suitable for use as control antibody for CYP24 siRNA (h): sc-44652, CYP24 siRNA (m): sc-44653, CYP24 shRNA Plasmid (h): sc-44652-SH, CYP24 shRNA Plasmid (m): sc-44653-SH, CYP24 shRNA (h) Lentiviral Particles: sc-44652-V and CYP24 shRNA (m) Lentiviral Particles: sc-44653-V.


Molecular Weight of CYP24: 59 kDa.

Positive Controls: mouse liver extract: sc-2256, MIA PaCa-2 cell lysate: sc-2285 or mouse pancreas tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CYP24 (C-18): sc-32166. Western blot analysis of CYP24 expression in mouse pancreas (A) and mouse liver (B) tissue extracts.

SELECT PRODUCT CITATIONS

- Lopes, N., et al. 2010. Alterations in vitamin D signalling and metabolic pathways in breast cancer progression: a study of VDR, CYP27B1 and CYP24A1 expression in benign and malignant breast lesions. BMC Cancer 10: 483.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **CYP24 (E-7): sc-365700**, our highly recommended monoclonal alternative to CYP24 (C-18).