

Coronin 1B (S-20): sc-32214

BACKGROUND

Coronins are a family of WD repeat-containing, actin-binding proteins that localize to submembrane areas and regulate cell motility and cytoskeletal rearrangement. Coronin 1A (CORO1A, CLIPINA, CLABP, TACO, p57) can form coiled-coil-mediated homotrimeric complexes that influence early phagosome formation. PKC-dependent phosphorylation of Coronin 1B (CORO1B) at serine 2 regulates leading edge dynamics and cell motility in fibroblasts through interactions with Arp2/3 complex. Coronin 1C (CORO1C, Coronin 3, HCRNN4) is abundant in differentiating Neuro-2a cells, PC-12 cells and primary oligodendrocytes, where it is thought to influence neuron morphogenesis and migration. Coronin 2A (CORO2A, CLIPINB, IR10, WDR2) is a component of the approximately 1.5-2 megadalton N-CoR (nuclear receptor corepressor) complex of 10-12 proteins, which recruits HDACs to generate repressive chromatin. Coronin 7 (CORO7, CRN7) localizes to the Golgi membrane and influences the organization of intracellular membrane compartments and vesicular trafficking. Coronin 2B (CORO2B, CLIPINC) and Coronin 6 (CORO6) are similar to other members of this family, since they possess a conserved basic N-terminal motif and 3-10 WD repeats clustered in one to two core domains.

CHROMOSOMAL LOCATION

Genetic locus: CORO1B (human) mapping to 11q13.2; Coro1b (mouse) mapping to 19 A.

SOURCE

Coronin 1B (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Coronin 1B of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32214 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Coronin 1B (S-20) is recommended for detection of Coronin 1B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Coronin 1B (S-20) is also recommended for detection of Coronin 1B in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Coronin 1B siRNA (h): sc-44695, Coronin 1B siRNA (m): sc-44696, Coronin 1B shRNA Plasmid (h): sc-44695-SH, Coronin 1B shRNA Plasmid (m): sc-44696-SH, Coronin 1B shRNA (h) Lentiviral Particles: sc-44695-V and Coronin 1B shRNA (m) Lentiviral Particles: sc-44696-V.


Molecular Weight of Coronin 1B: 57 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, WI-38 whole cell lysate: sc-364260 or RAT2 whole cell lysate: sc-364198.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Coronin 1B (S-20): sc-32214. Western blot analysis of Coronin 1B expression in NIH/3T3 (A), WI-38 (B) and RAT2 (C) whole cell lysates.



Coronin 1B (S-20): sc-32214. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Coronin 1B (G-8): sc-271445** or **Coronin 1B (E-3): sc-271375**, our highly recommended monoclonal alternatives to Coronin 1B (S-20).