

SOD-3 (N-19): sc-32219

BACKGROUND

The superoxide dismutase family is composed of three metalloenzymes (SOD-1, SOD-2 and SOD-3) that catalyze the oxido-reduction of reactive oxygen species (ROS) such as superoxide anion. ROS are implicated in a wide range of degenerative processes, including Alzheimer disease, Parkinson disease and ischemic heart disease. Cu/Zn superoxide dismutase-1 (SOD-1) is a well characterized cytosolic scavenger of oxygen free radicals that requires copper and zinc binding to potentiate its enzymatic activity. The SOD-2 precursor is a 222 amino acid protein that is encoded by nuclear chromatin, synthesized in the cytosol and imported posttranslationally into the mitochondrial matrix. SOD-3, also designated extracellular superoxide dismutase (EC-SOD), is an extracellular zinc and copper binding protein that destroys radicals that are toxic to biological systems but that are normally produced within cells. SOD-3 is found in extracellular fluids such as lymph, plasma and synovial fluid.

REFERENCES

- Levanon, D., et al. 1985. Architecture and anatomy of the chromosomal locus in human chromosome 21 encoding the Cu/Zn superoxide dismutase. *EMBO J.* 4: 77-84.
- Bewley, G.C. 1988. cDNA and deduced amino acid sequence of murine Cu/Zn superoxide dismutase. *Nucleic Acids Res.* 16: 2728.
- Beckman, J.S., et al. 1993. ALS, SOD and peroxynitrite. *Nature* 364: 584.
- Sandstrom, J., et al. 1994. 10-fold increase in human plasma extracellular superoxide dismutase content caused by a mutation in heparin-binding domain. *J. Biol. Chem.* 269: 19163-19166.
- Li, Y., et al. 1995. Dilated cardiomyopathy and neonatal lethality in mutant mice lacking manganese superoxide dismutase. *Nat. Genet.* 11: 376-381.
- Adachi, T., et al. 1996. An Arginine 213 to glycine mutation in human extracellular-superoxide dismutase reduces susceptibility to trypsin-like proteinases. *J. Biochem.* 120: 184-188.

CHROMOSOMAL LOCATION

Genetic locus: SOD3 (human) mapping to 4p15.2; Sod3 (mouse) mapping to 5 C1.

SOURCE

SOD-3 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SOD-3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32219 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SOD-3 (N-19) is recommended for detection of SOD-3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SOD-3 siRNA (h): sc-44699, SOD-3 shRNA Plasmid (h): sc-44699-SH and SOD-3 shRNA (h) Lentiviral Particles: sc-44699-V.


Molecular Weight of SOD-3: 32 kDa.

Positive Controls: Daudi cell lysate: sc-2415.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SOD-3 (N-19): sc-32219. Western blot analysis of SOD-3 expression in JAR (A) and WI 38 (B) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **SOD-3 (G-11): sc-376948** or **SOD-3 (4G11G6): sc-101338**, our highly recommended monoclonal alternatives to SOD-3 (N-19).