

Ly-GDI (T-18): sc-32226

BACKGROUND

The Ras superfamily of small GTP-binding proteins are critical mediators of diverse cell signaling pathways, including those leading to proliferation, cytoskeletal organization and secretion. The counter-conversion of the active GTP-bound form of these proteins to their inactive GDP-bound form is influenced by two types of regulatory proteins: those that alter the intrinsic GTPase activity of the GTP-binding proteins and those that alter the rate of GDP/GTP exchange. Guanine nucleotide-releasing factors (GRFs) increase the GDP dissociation rate, while GDP-dissociation inhibitors (GDIs) decrease the dissociation rate. The Rho GDI subfamily is composed of Rho GDI α , Ly-GDI (also known as Rho GDI β and previously known as GDI/D4) and Rho GDI γ . The Rho GDI proteins interact with and have varying affinities for several Ras-like GTP binding proteins, including Rho A, Rho B, Rac and Cdc42. Ly-GDI is expressed only in hematopoietic cells, predominantly in B and T lymphocyte cell lines.

REFERENCES

1. Trahey, M., et al. 1987. A cytoplasmic protein stimulates normal N-Ras p21 GTPase, but does not affect oncogenic mutants. *Science* 238: 542-545.
2. Hall, A. 1990. The cellular functions of small GTP-binding proteins. *Science* 249: 635-640.
3. Bourne, H.R., et al. 1990. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348: 125-132.
4. Garrett, M.D., et al. 1991. Purification and N-terminal sequence of the p21Rho GTPase-activating protein, Rho GAP. *Biochem. J.* 276: 833-836.
5. Scherle, P., et al. 1993. Ly-GDI, a GDP-dissociation inhibitor of the RhoA GTP-binding protein, is expressed preferentially in lymphocytes. *Proc. Natl. Acad. Sci. USA* 90: 7568-7572.
6. Platko, J.V., et al. 1995. A single residue can modify target-binding affinity and activity of the functional domain of the Rho-subfamily GDP dissociation inhibitors. *Proc. Natl. Acad. Sci. USA* 92: 2974-2978.

CHROMOSOMAL LOCATION

Genetic locus: ARHGDI β (human) mapping to 12p12.3; Arhgdib (mouse) mapping to 6 G1.

SOURCE

Ly-GDI (T-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Ly-GDI of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32226 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Ly-GDI (T-18) is recommended for detection of Ly-GDI of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Ly-GDI siRNA (h): sc-35826, Ly-GDI siRNA (m): sc-35827, Ly-GDI shRNA Plasmid (h): sc-35826-SH, Ly-GDI shRNA Plasmid (m): sc-35827-SH, Ly-GDI shRNA (h) Lentiviral Particles: sc-35826-V and Ly-GDI shRNA (m) Lentiviral Particles: sc-35827-V.


Molecular Weight of Ly-GDI: 27 kDa.

Positive Controls: BJAB whole cell lysate: sc-2207, U-937 cell lysate: sc-2239 or HEL 92.1.7 cell lysate: sc-2270.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Ly-GDI (T-18): sc-32226. Western blot analysis of Ly-GDI expression in ALL-SIL (A), BJAB (B), HEL 92.1.7 (C) and U-937 (D) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **Ly-GDI (G-12): sc-376473** or **Ly-GDI (D-7): sc-271108**, our highly recommended monoclonal alternatives to Ly-GDI (T-18). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Ly-GDI (G-12): sc-376473**.