

RFRP-1 (S-16): sc-32374

BACKGROUND

The human RFamide-related peptide gene, RFRP (also designated NPVF or C7orf9), is responsible for encoding three small neuropeptides designated RFRP-1 (NPSF), RFRP-2 and RFRP-3 (NPVF). The homologous gene in rodents encodes only two functional neuropeptide: RFRP-1 (NPSF) and RFRP-3 (NPVF). RFamide-related peptides constitute a large family of neuropeptides in a wide range of species that are known to play a role in neurotransmission, neuro-modulation, cardioexcitation and control of muscle contraction. Neuropeptides RFRP-1 and RFRP-3 efficiently inhibit Forskolin-induced production of cAMP. RFRP-2, however, does not appear to have a similar inhibitory activity. RFamide-related peptides are secreted and abundantly expressed in retina. RFRP-1 and RFRP-3 are also widely distributed in fetal and adult brain, including the forebrain, hypothalamus, thalamus, midbrain, pons and medulla oblongata. RFRP-1 and the prolactin (PRL)-releasing peptide-31 (PrRP-31) may be involved in the stimulation of stress hormone secretion by either direct pituitary or indirect hypothalamic actions.

REFERENCES

- Hinuma, S., Shintani, Y., Fukusumi, S., Iijima, N., Matsumoto, Y., Hosoya, M., Fujii, R., Watanabe, T., Kikuchi, K., Terao, Y., Yano, T., Yamamoto, T., Kawamata, Y., Habata, Y., Asada, M., Kitada, C., Kurokawa, T., Onda, H., Nishimura, O., Tanaka, M., Ibata, Y. and Fujino, M. 2000. New neuropeptides containing carboxy-terminal RFamide and their receptor in mammals. *Nat. Cell Biol.* 2: 703-708.
- Fukusumi, S., Habata, Y., Yoshida, H., Iijima, N., Kawamata, Y., Hosoya, M., Fujii, R., Hinuma, S., Kitada, C., Shintani, Y., Suenaga, M., Onda, H., Nishimura, O., Tanaka, M., Ibata, Y. and Fujino, M. 2001. Characteristics and distribution of endogenous RFamide-related peptide-1. *Biochim. Biophys. Acta* 1540: 221-232.
- Schulz, H.L., Stoehr, H., White, K., van Driel, M.A., Hoyng, C.B., Cremers, F. and Weber, B.H. 2002. Genomic structure and assessment of the retinally expressed RFamide-related peptide gene in dominant cystoid macular dystrophy. *Mol. Vis.* 8: 67-71.
- Yano, T., Iijima, N., Kakihara, K., Hinuma, S., Tanaka, M. and Ibata, Y. 2003. Localization and neuronal response of RFamide related peptides in the rat central nervous system. *Brain Res.* 982: 156-167.
- Samson, W.K., Keown, C., Samson, C.K., Samson, H.W., Lane, B., Baker, J.R. and Taylor, M.M. 2003. Prolactin-releasing peptide and its homolog RFRP-1 act in hypothalamus but not in anterior pituitary gland to stimulate stress hormone secretion. *Endocrine* 20: 59-66.
- Yano, T., Iijima, N., Hinuma, S., Tanaka, M. and Ibata, Y. 2004. Developmental expression of RFamide-related peptides in the rat central nervous system. *Brain Res. Dev. Brain Res.* 152: 109-120.
- Johnson, M.A., Tsutsui, K. and Fraley, G.S. 2007. Rat RFamide-related peptide-3 stimulates GH secretion, inhibits LH secretion, and has variable effects on sex behavior in the adult male rat. *Horm. Behav.* 51: 171-180.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: RFRP (human) mapping to 7p15.3.

SOURCE

RFRP-1 (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of RFamide-related peptides precursor protein of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32374 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

RFRP-1 (S-16) is recommended for detection of RFamide-related peptides precursor protein and RFRP-1 active peptide of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RFRP-1 (S-16) is also recommended for detection of RFamide-related peptides precursor protein and RFRP-1 active peptide in additional species, including porcine.

Suitable for use as control antibody for RFRP siRNA (h): sc-44797, RFRP shRNA Plasmid (h): sc-44797-SH and RFRP shRNA (h) Lentiviral Particles: sc-44797-V.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.