

# CMAH (V-18): sc-323744

## BACKGROUND

The sialic acids are a family of acidic sugars typically found in the outer portion of the cell surface and in secreted glycoconjugates of all vertebrates. Cell membrane sialic acid is involved in cell-cell and cell-pathogen interactions and in binding of cells to the extracellular matrix. The two most common forms of sialic acid found in mammalian cells are N-acetylneuraminic acid (Neu5Ac) and its hydroxylated derivative, N-glycolylneuraminic acid (Neu5Gc). CMAH (cytidine monophospho-N-acetylneuraminic acid hydroxylase), also known as CMP-Neu5Ac hydroxylase or CMP-N-acetylneuraminic acid monooxygenase, is a 577 amino acid cytoplasmic protein that is expressed in all tissues, except in brain. Belonging to the CMP-Neu5Ac hydroxylase family, CMAH catalyzes the conversion of CMP-Neu5Ac into its hydroxylated derivative CMP-Neu5Gc, a sialic acid abundantly expressed at the surface of many cells. CMAH exists as two isoforms due to alternative splicing events. Isoform 2 is expressed in the endoplasmic reticulum.

## REFERENCES

1. Kawano, T., et al. 1995. Molecular cloning of cytidine monophospho-N-acetylneuraminic acid hydroxylase. Regulation of species- and tissue-specific expression of N-glycolylneuraminic acid. *J. Biol. Chem.* 270: 16458-16463.
2. Muchmore, E.A., et al. 1998. A structural difference between the cell surfaces of humans and the great apes. *Am. J. Phys. Anthropol.* 107: 187-198.
3. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 603209. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Chou, H.H., et al. 2002. Inactivation of CMP-N-acetylneuraminic acid hydroxylase occurred prior to brain expansion during human evolution. *Proc. Natl. Acad. Sci. USA* 99: 11736-11741.
5. Bighignoli, B., et al. 2007. Cytidine monophospho-N-acetylneuraminic acid hydroxylase (CMAH) mutations associated with the domestic cat AB blood group. *BMC Genet.* 8: 27.
6. Naito, Y., et al. 2007. Germinal center marker GL7 probes activation-dependent repression of N-glycolylneuraminic acid, a sialic acid species involved in the negative modulation of B-cell activation. *Mol. Cell. Biol.* 27: 3008-3022.
7. Hedlund, M., et al. 2007. N-glycolylneuraminic acid deficiency in mice: implications for human biology and evolution. *Mol. Cell. Biol.* 27: 4340-4346.
8. Varki, A. 2009. Multiple changes in sialic acid biology during human evolution. *Glycoconj. J.* 26: 231-245.
9. Rich, S.M., et al. 2009. The origin of malignant malaria. *Proc. Natl. Acad. Sci. USA* 106: 14902-14907.

## CHROMOSOMAL LOCATION

Genetic locus: CMAHP (human) mapping to 6p22.3.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## SOURCE

CMAH (V-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CMAH of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-323744 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

CMAH (V-18) is recommended for detection of CMAH of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with CMAS.

CMAH (V-18) is also recommended for detection of CMAH in additional species, including canine and bovine.

Molecular Weight of CMAH: 8 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **CMAH (E-7): sc-365023**, our highly recommended monoclonal alternative to CMAH (V-18).