

CTAGE5 (E-16): sc-323755

BACKGROUND

Cutaneous T cell lymphomas (CTCL) represent a group of malignancies that originate from CD4-T lymphocytes and manifest on the skin. CTCL is a general term for several neoplasms including mycosis fungoides, T cell leukemia/lymphoma and pagetoid reticulosis, all of which are very difficult to treat in the advanced stages. CTAGE5 (cutaneous T-cell lymphoma-associated antigen 5), also known as MEA11, MEA6, MGEA11 or MGEA6, is an 804 amino acid tumor associated antigen that is found in tumors of various origins, including cutaneous T cell lymphomas. CTAGE5 is expressed as multiple isoforms due to alternative splicing events. Isoform 5A is expressed only in testis at the protein level while other isoforms are expressed in several other normal tissues, including brain, muscle and cranial skin.

REFERENCES

1. Eichmüller, S., et al. 2001. Serological detection of cutaneous T cell lymphoma-associated antigens. *Proc. Natl. Acad. Sci. USA* 98: 629-634.
2. Comtesse, N., et al. 2001. The MGEA6 multigene family has an active locus on 14q and at least nine pseudogenes on different chromosomes. *Genomics* 75: 43-48.
3. Comtesse, N., et al. 2002. MGEA6 is tumor-specific overexpressed and frequently recognized by patient-serum antibodies. *Oncogene* 21: 239-247.
4. Eichmüller, S. 2002. Towards defining specific antigens for cutaneous lymphomas. *Onkologie* 25: 448-454.
5. Eichmüller, S., et al. 2003. Tumor-specific antigens in cutaneous T cell lymphoma: expression and sero-reactivity. *Int. J. Cancer* 104: 482-487.
6. Usener, D., et al. 2003. cTAGE: a cutaneous T cell lymphoma associated antigen family with tumor-specific splicing. *J. Invest. Dermatol.* 121: 198-206.
7. Atanackovic, D., et al. 2006. Expression of cancer-testis antigens as possible targets for antigen-specific immunotherapy in head and neck squamous cell carcinoma. *Cancer Biol. Ther.* 5: 1218-1225.

CHROMOSOMAL LOCATION

Genetic locus: CTAGE5 (human) mapping to 14q21.1; Ctage5 (mouse) mapping to 12 C1.

SOURCE

CTAGE5 (E-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CTAGE5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-323755 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CTAGE5 (E-16) is recommended for detection of CTAGE5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other CTAGE family members.

CTAGE5 (E-16) is also recommended for detection of CTAGE5 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for CTAGE5 siRNA (h): sc-92131, CTAGE5 siRNA (m): sc-142612, CTAGE5 shRNA Plasmid (h): sc-92131-SH, CTAGE5 shRNA Plasmid (m): sc-142612-SH, CTAGE5 shRNA (h) Lentiviral Particles: sc-92131-V and CTAGE5 shRNA (m) Lentiviral Particles: sc-142612-V.


Molecular Weight of CTAGE5: 91 kDa.

Positive Controls: PC-3 cell lysate: sc-2220.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CTAGE5 (E-16): sc-323755. Western blot analysis of CTAGE5 expression in PC-3 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

MONOS
Satisfaction
Guaranteed

Try **CTAGE2/5 (B-1): sc-365330**, our highly recommended monoclonal alternative to CTAGE5 (E-16).