HSPA7 (L-13): sc-323963



The Power to Question

BACKGROUND

The heat shock proteins (HSPs) comprise a group of highly conserved, abundantly expressed proteins with diverse functions, including the assembly and sequestering of multi-protein complexes, the transportation of nascent polypeptide chains across cellular membranes and the regulation of protein folding. HSPA7 (heat shock 70 kDa protein 7), also known as HSP70B, is a 367 amino acid protein belonging to the heat shock protein 70 family, and like other HSP proteins, mediates protein folding within the cytosol, as well as within other organelles throughout the cell. The gene encoding HSPA7 maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

- Voellmy, R., et al. 1985. Isolation and functional analysis of a human 70,000-dalton heat shock protein gene segment. Proc. Natl. Acad. Sci. USA 82: 4949-4953.
- 2. Schiller, P., et al. 1988. *Cis*-acting elements involved in the regulated expression of a human HSP70 gene. J. Mol. Biol. 203: 97-105.
- 3. Leung, T.K., et al. 1990. The human heat-shock protein family. Expression of a novel heat-inducible HSP70 (HSP70B') and isolation of its cDNA and genomic DNA. Biochem. J. 267: 125-132.
- 4. Leung, T.K., et al. 1992. The human heat-shock genes HSPA6 and HSPA7 are both expressed and localize to chromosome 1. Genomics 12: 74-79.
- 5. Grosz, M.D., et al. 1992. Syntenic conservation of HSP70 genes in cattle and humans. Genomics 14: 863-868.
- Brzustowicz, L.M., et al. 2002. Fine mapping of the schizophrenia susceptibility locus on chromosome 1q22. Hum. Hered. 54: 199-209.
- 7. Online Mendelian Inheritance in Man, OMIM™. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 140555. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Liu, Y., et al. 2008. HSP70 is associated with endothelial activation in placental vascular diseases. Mol. Med. 14: 561-566.

CHROMOSOMAL LOCATION

Genetic locus: HSPA7 (human) mapping to 1g23.3.

SOURCE

HSPA7 (L-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of HSPA7 of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-323963 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HSPA7 (L-13) is recommended for detection of HSPA7 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other HSPA family members.

Molecular Weight of HSPA7: 40 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**