

MYH3 (N-16): sc-324153



The Power to Question

BACKGROUND

Actin is a highly conserved protein that is expressed in all eukaryotic cells. Actin filaments can form both stable and labile structures, and are crucial components of microvilli and the contractile apparatus of muscle cells. Myosin is a hexamer of two heavy chains (MHC) and four light chains (MLC) that interact with actin to generate the force for diverse cellular movements, including cytokinesis, phagocytosis and muscle contraction. MYH3 (Myosin heavy chain 3), also known as muscle embryonic Myosin heavy chain or SMHCE, is a 1,940 amino acid that localizes to the thick filaments of myofibrils. While highly expressed in fetal skeletal muscle, MYH3 is barely detectable in adult skeletal muscle. Defects in the gene encoding MYH3, which maps to human chromosome 17p13.1, are the cause of distal arthrogyrosis type 2B (DA2B).

REFERENCES

1. Nagai, R., Kuro-o, M., Babij, P. and Periasamy, M. 1989. Identification of two types of smooth muscle Myosin heavy chain isoforms by cDNA cloning and immunoblot analysis. *J. Biol. Chem.* 264: 9734-9737.
2. Karsch-Mizrachi, I., Feghali, R., Shows, T.B. and Leinwand, L.A. 1990. Generation of a full-length human perinatal Myosin heavy-chain-encoding cDNA. *Gene* 89: 289-294.
3. Bober, E., Buchberger-Seidl, A., Braun, T., Singh, S., Goedde, H. W. and Arnold, H.H. 1990. Identification of three developmentally controlled isoforms of human Myosin heavy chains. *Eur. J. Biochem.* 189: 55-65.
4. Hughes, S.M. and Blau H.M. 1992. Muscle fiber pattern is independent of cell lineage in postnatal rodent development. *Cell* 68: 659-671.
5. Cheney, R.E., Riley, M.A. and Mooseker, M.S. 1993. Phylogenetic analysis of the Myosin superfamily. *Cell Motil. Cytoskeleton.* 24: 215-223.
6. Jullian, E.H., Kelly, A.M., Pompidou, A.J., Hoffman, R., Schiaffino, S., Stedman, H.H. and Rubinstein, N.A. 1995. Characterization of a human perinatal Myosin heavy-chain transcript. *Eur. J. Biochem.* 230: 1001-1006.
7. Owens, G.K. 1995. Regulation of differentiation of vascular smooth muscle cells. *Physiol. Rev.* 75: 487-517.
8. Weiss, A. and Leinwand, L.A. 1996. The mammalian Myosin heavy chain gene family. *Annu. Rev. Cell Dev. Biol.* 12: 417-439.

CHROMOSOMAL LOCATION

Genetic locus: MYH3 (human) mapping to 17p13.1; Myh3 (mouse) mapping to 11 B3.

SOURCE

MYH3 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of MYH3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-324153 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MYH3 (N-16) is recommended for detection of MYH3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other MYH family members.

MYH3 (N-16) is also recommended for detection of MYH3 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for MYH3 siRNA (h): sc-93798, MYH3 siRNA (m): sc-149742, MYH3 shRNA Plasmid (h): sc-93798-SH, MYH3 shRNA Plasmid (m): sc-149742-SH, MYH3 shRNA (h) Lentiviral Particles: sc-93798-V and MYH3 shRNA (m) Lentiviral Particles: sc-149742-V.

Molecular Weight of MYH3: 224 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **MYH3 (F1.652): sc-53091** or **MYH (B-5): sc-376157**, our highly recommended monoclonal alternatives to MYH3 (N-16).