



## MSL-2 (dC-20): sc-32459

### BACKGROUND

*Drosophila melanogaster* is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes can be categorized based on the type of protein for which they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing and chromatin associated) or other functional proteins. The male-specific lethal 2 (MSL-2) protein is essential for X-chromosome dosage compensation. Female flies inhibit MSL-2 mRNA translation, which is mediated by sex-lethal (SXL) in both the 5' and 3' untranslated regions.

### REFERENCES

1. Adams, M.D., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
2. Forch, P., et al. 2001. Modulation of MSL-2 5' splice site recognition by sex-lethal. *RNA* 7: 1185-1191.
3. Gebauer, F., et al. 2003. *Drosophila* sex-lethal inhibits the stable association of the 40S ribosomal subunit with MSL-2 mRNA. *Mol. Cell*. 11: 1397-1404.
4. Grskovic, M., et al. 2003. A co-repressor assembly nucleated by sex-lethal in the 3'UTR mediates translational control of *Drosophila* MSL-2 mRNA. *EMBO J.* 22: 5571-5581.
5. Beckmann, K. et al. 2005. A dual inhibitory mechanism restricts MSL-2 mRNA translation for dosage compensation in *Drosophila*. *Cell* 122: 529-540.
6. The Interactive Fly. <http://sdb.bio.purdue.edu/fly/aimain/1aahome.htm>.  
<http://sdb.bio.purdue.edu/fly/aimain/6biochem.htm>.

### SOURCE

MSL-2 (dC-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of MSL-2 of *Drosophila melanogaster* origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32459 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### APPLICATIONS

MSL-2 (dC-20) is recommended for detection of MSL-2 of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of MSL-2: 97 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.